

(12)

EUROPEAN PATENT SPECIFICATION

(54) Date of publication of patent specification: **04.04.90**

(51) Application number: **84201812.9**

(72) Date of filing: **06.12.84**

(51) Int. Cl.⁵: **C 07 D 471/04,**
C 07 D 401/12,
C 07 D 235/30, A 61 K 31/445

(54) **N-(4-piperidinyl) bicyclic condensed 2-imidazolamine derivatives.**

(10) Priority: **09.01.84 US 569115**
15.10.84 US 660670

(43) Date of publication of application:
21.08.85 Bulletin 85/34

(53) Publication of the grant of the patent:
04.04.90 Bulletin 90/14

(54) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

(56) References cited:
AT-B-315 831
GB-A-1 260 857
US-A-2 971 005
US-A-4 219 559

CHEMICAL ABSTRACTS, vol. 96, no. 23, 07 June 1982, Columbus, OH (US); P.M.LADURON: "In vitro and in vivo binding characteristics of a new long-acting histamine H1 antagonist astemizole", p. 62, no. 193269h

CHEMICAL ABSTRACTS, vol. 100, no. 17, 23 April 1984, Columbus, OH (US); M.FABA: "1-(4-

The file contains technical information submitted after the application was filed and not included in this specification

(71) Proprietor: **JANSSEN PHARMACEUTICA N.V.**
Turnhoutsebaan 30
B-2340 Beerse (BE)

(72) Inventor: **Janssens, Frans Eduard**
Tinstraat 79
B-2830 Bonheiden (BE)
 Inventor: **Torremans, Joseph Leo Ghislanus**
Lijsterstraat 11
B-2340 Beerse (BE)
 Inventor: **Hens, Jozef Francis**
Rector de Ramstraat 54
B-2260 Nijlen (BE)
 Inventor: **Van Offenwert, Theophilus Theresia**
JoannesMaria
Kardinaal Cardijnlaan 53
B-2250 Vosselaar (BE)

(56) References cited:
fluorophenyl)methyl-2-1-(4-methoxyphenyl)-4-piperidinylaminorbenzimidazole", p. 652, no. 139109u


CHEMICAL ABSTRACTS, vol. 100, no. 1, 02 Jan 1984, Columbus, OH (US); J.B.A.THUJSEN: "Synthesis of tritium- and carbon-14-labeled astemizole", p. 552, no. 6394m

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European patent convention).

Description

Background of the invention

In U.S. Patent No. 4,219,559 there are described a number of *N*-heterocyclyl-4-piperidinamines having the formula




which compounds are useful as antihistaminic agents.

The compounds of the present invention differ from the prior art compounds essentially by the nature of the 1-piperidinyl substituent.


Description of the preferred embodiments

This invention is concerned with novel *N*-heterocyclyl-4-piperidinamines which may structurally be represented by the formula



the pharmaceutically acceptable acid addition salts and the possible stereochemically isomeric forms thereof, wherein:

$A^1=A^2=A^3=A^4$ is a bivalent radical having the formula




wherein one or two hydrogen atoms in said radicals (a-1) — (a-5) may, each independently from each other, be replaced by halo, lower alkyl, lower alkyloxy, trifluoromethyl or hydroxy;


R is a member selected from the group consisting of hydrogen and lower alkyl;

R¹ is a member selected from the group consisting of hydrogen, alkyl, cycloalkyl, Ar¹ and lower alkyl substituted with one or two Ar¹ radicals;

R² is a member selected from the group consisting of hydrogen, lower alkyl, cycloalkyl, (lower alkyl)-CO—, (lower alkyl)OCO— and Ar²-lower alkyl; and

L is a radical of formula





(b-4);

Het-Alk-

(b-5);

(lower alkenyl)-Y¹-Alk-


(b-6); or

i) where A¹=A²=A³=A⁴ is a radical of formula (a-3), (a-4) or (a-5), or

ii) where A¹=A²=A³=A⁴ is a radical of formula (a-1) or (a-2), and R¹ is Ar³ or lower alkyl substituted with one or two Ar² radicals, said Ar² being pyrazinyl, thiazolyl or imidazolyl, optionally substituted with lower alkyl; L may also be a radical of formula Ar¹-Alk- (b-7);


said W being a member selected from the group consisting of hydrogen, lower alkyl, Ar¹, Ar¹-lower alkyl, 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl,

a radical of formula



(c-1-a),

a radical of formula




(c-1-b), or

a radical of formula W¹-Z¹- (c-1-c), wherein R³ and R⁴ are each independently hydrogen or lower alkyl; and W¹ is cycloalkyl or lower alkyl, optionally substituted with up to two substituents selected from the group consisting of hydroxy, lower alkyloxy, 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl and Ar¹; and where Z¹ is NR⁶, W¹ may also be hydrogen, amino, lower alkylamino, Ar¹-amino or nitro;


said W² being a member selected from the group consisting of hydrogen, lower alkyl, Ar¹ and a radical of formula R⁵-Z¹- (c-2-a), wherein R⁵ is hydrogen, lower alkyl or Ar¹;

said T being a radical of formula




R⁷-SO₂-NR⁸- (c-3-b); R⁶ being hydrogen, lower alkyl or Ar¹; R⁷ being lower alkyl or Ar¹; and R⁸ being hydrogen or lower alkyl;

said Het being a radical of formula (c-1-a), (c-1-b), or a radical of formula



or of formula




or a radical R¹³ of formula

(c-4-c),

wherein R⁹, R¹⁰, R¹¹ and R¹² are each independently hydrogen or lower alkyl; and wherein R¹³ is hydrogen, lower alkyl or amino; or

said Het being furan substituted with lower alkyl, said lower alkyl being optionally substituted with hydroxy, mercapto, lower alkyloxy, lower alkylthio, (aminolower alkyl)thio, Ar¹-O- or with a radical of



(c-4-d-1),

s being an integer of from 1 to 6 inclusive; or where Z or Y is a direct bond, s may also be 0; and R¹⁴ being hydrogen or lower alkyl;

wherein: n is 0 or the integer 1 or 2;

X is O, S, NR¹⁵ or CHNO₂;

Y is O, S, NR¹⁶ or a direct bond;

Y¹ is O, S or NR¹⁶;

Y² is S or NR¹⁶;

Z is O, S, NR⁸ or a direct bond;

Z¹ is O, S or NR⁸;

X² and Y³ independently having the same meaning of X respectively Y;

said R¹⁵ being hydrogen, lower alkyl, cyano, nitro, Ar²-sulfonyl, lower alkylsulfonyl, lower alkylcarbonyl or Ar²-carbonyl;

said R¹⁶ being hydrogen, lower alkyl, (Ar²)lower alkyl, 2-lower alkoxy-1,2-dioxoethyl; or a radical of formula —C(=X)—R^{17} ;

R¹⁷ being hydrogen, lower alkyl, Ar², Ar²-lower alkyl, lower alkoxy, Ar²-lower alkoxy, mono- or di(lower alkyl)amino, Ar²-lower alkylamino or Ar²-lower alkyl(lower alkyl)amino;

wherein Ar¹ is a member selected from the group consisting of phenyl, being optionally substituted with up to three substituents each independently selected from the group consisting of halo, hydroxy, nitro, cyano, trifluoromethyl, lower alkyl, lower alkoxy, lower alkylthio, mercapto, amino, mono- and di(lower alkyl)amino, carboxyl, lower alkylcarbonyl and lower alkyl-CO—; thienyl; halothienyl; furanyl; lower alkyl substituted furanyl; pyridinyl; pyrazinyl; thiazonyl and imidazolyl optionally substituted with lower alkyl; and wherein Ar² is a member selected from the group consisting of phenyl being optionally

substituted with up to three substituents each independently selected from the group consisting of halo, hydroxy, nitro, cyano, trifluoromethyl, lower alkyl, lower alkoxy, lower alkylthio, mercapto, amino, mono- and di(lower alkyl)amino, carboxyl, lower alkylcarbonyl and (lower alkyl)-CO;

provided that:

i) when A¹=A²=A³=A⁴ is a radical of formula (a-1) or (a-2), and L is a radical of formula (b-1), wherein W is other than hydrogen or other than a radical of formula (c-1-a) or (c-1-b), then X is other than O;

ii) when L is a radical of formula (b-1), wherein W is a radical of formula (c-1-c), wherein Z¹ is NH then W¹ is other than hydrogen or lower alkyl;

iii) when A¹=A²=A³=A⁴ is a radical of formula (a-1) or (a-2), and L is a radical of formula (b-3), wherein X is O, Y is NR¹⁶, O or a direct bond, and X² is O,

a) then Y³ is not 0;

b) and W² being lower alkyl then Y³ is not a direct bond;

As used in the foregoing definitions the term halo is generic to fluoro, chloro, bromo and iodo; the term "lower alkyl" is meant to include straight- and branched-chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as, for example, methyl, ethyl, 1-methylethyl, 1,1-dimethylethyl, propyl, 2-methylpropyl, butyl, pentyl, hexyl; "alkyl" is meant to include lower alkyl radicals, as defined hereinabove, and the higher homologs thereof having from 7 to 10 carbon atoms; the term "cycloalkyl" is generic to cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; "lower alkenyl" is meant to include straight- or

branched-chain hydrocarbon radicals containing one double bond, and having from 2 to 6 carbon atoms such as, for example, ethenyl, 2-propenyl, 2-butenyl, 3-pentenyl and 3-hexenyl; and "lower alkanediyl" is meant to include bivalent straight- or branched-chain alkanediyl radicals having from 1 to 6 carbon atoms.

Some of the compounds of formula (I) may contain in their structure a keto-enol tautomeric system or a vinyllog system thereof and consequently these compounds may be present in their keto form as well as their enol form.

Preferred compounds within the invention are those wherein:

i) L is a radical of formula (b-1), wherein Y is NH, X is O and W is hydrogen; or L is a radical of formula (b-1) wherein X is S, NH or CN, Y is NH and W is 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl, or a radical of formula (c-1-c), wherein Z¹ is NR⁸ and W¹ is amino, nitro or lower alkyl, optionally substituted with one hydroxy, lower alkoxy, 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl or phenyl radical, or with two lower alkoxy radicals; or L is a radical of formula (b-1), wherein X is S, NH or CN, Y is NH and W is lower alkoxy of lower alkylthio; or wherein L is a radical of formula (b-1) wherein W is a radical of formula (c-1-a) or (c-1-b);

ii) L is a radical of formula (b-2) wherein n is 1, X is O or S and W is a radical of formula (c-1-c), wherein Z¹ is NR⁸ and W¹ is lower alkyl;


iii) L is a radical of formula (b-3), wherein X is O, Y is NH, X² is O, YH³ is NR¹⁶ and W² is lower alkyl;

iv) L is a radical of formula (b-4), wherein T is a radical of formula (c-3-a), wherein X is O or S, Z is NR⁸ and R² is hydrogen or lower alkyl; or wherein T is a radical of formula (c-3-b), wherein R² is hydrogen and R⁷ is lower alkyl;

v) L is a radical of formula (b-5) wherein Het is a radical of formula (c-4-a), wherein R⁹, R¹¹ and R¹² are hydrogen; or wherein Het is a radical of formula (c-4-c); or wherein Het is furan substituted with lower alkyl being substituted with hydroxy or with a radical of formula (c-4-d-1), wherein Y is O or S, Z is NH or a direct bond and R¹⁴ is hydrogen;


vi) L is a radical of formula (b-6) wherein Y¹ is O;

vii) L is a radical of formula (b-7) wherein Ar¹ is phenyl substituted with hydroxy or lower alkyloxy.
In order to simplify the structural representations of the compounds of formula (I) and of certain precursors and intermediates thereof, the




represented by the symbol D .

The compounds of formula (I) can generally be prepared by reacting an intermediate of formula (II) with a piperidine derivative of formula (III), following art-known alkylating procedures.







Q¹ and Q² are selected so that during the alkylation reaction a radical or formula L is formed.

For example, the compounds of formula (I) can generally be prepared by N-alkylating a piperidine of formula (III) wherein Q² is hydrogen, said piperidine being represented by the formula (III-a), with a reagent of formula (II) wherein Q¹ has the general formula L—G, (II-a).



In (II-a) G represents an appropriate reactive leaving group such as, for example, halo, e.g., chloro, bromo or iodo, or a sulfonyloxy group, e.g. methylsulfonyloxy or 4-methylphenylsulfonyloxy.




Additionally, the compounds of formula (I) wherein L is a radical of formula (b-1) or (b-3) wherein Y is Y¹, or wherein L is a radical of formula (b-6) or (b-2), said compounds being represented by the formulae (I-a-1), respectively (I-a-2), (I-a-3) and (I-a-4) can be prepared by alkylating a piperidine of formula (III-b-1), respectively (III-b-2) with a reagent of formula (II-b-1) respectively (II-b-2), (II-b-3) and (II-b-4).




G¹ having the previously defined meaning of G and, where G¹ is connected to >C=X it may also represent a lower alkoxy, a lower alkylthio, an Ar²-oxy, an Ar²-thio, a lower alkylcarbonyloxy, or a lower

alkyloxycarbonyloxy group, and where G¹ is connected to >C=N—R¹⁸, it may also be —N(lower alkyl)NO.

The compounds of formula (I-a-1), (I-a-2), (I-a-3), and the compounds of formula (I), wherein L is a radical of formula (b-5), wherein Het is a radical of formula (c-4-a), (c-4-b) or (c-4-c), said Het being represented by Het' and said compounds being represented by the formula (I-a-5), may also be prepared by alkylating a piperidine of formula (III), wherein Q² is a radical of formula —Alk—G, said piperidine being represented by the formula (III-c), with a reagent of formula (II-c-1), respectively (III-c-2), (II-c-3) and (II-c-4).




The compounds of formula (I-a-1) or (I-a-4), wherein W is W'—Z'—, said compounds being represented by the formula (I-a-1-a), respectively (I-a-4-a), may also be prepared by reacting a reagent of formula (II-d) with an intermediate of formula (III-b-1) respectively (III-b-2) in the presence of an appropriate >C=X generating agent such as, for example, urea, thiourea, 1,1'-carbonylbis(1H-imidazole), lower alkylcarbonylhalide, carbonyl chloride, or thiocarbonyl chloride



The alkylation reactions are conveniently conducted in an inert organic solvent such as, for example, an aromatic hydrocarbon, e.g., benzene, methylbenzene or dimethylbenzene; a lower alcohol, e.g., methanol, ethanol or 1-butanol; a ketone, e.g., 2-propanone, or 4-methyl-2-pentanone; an ether, e.g., 1,4-dioxane, 1,1'-oxybisethane or tetrahydrofuran; *N,N*-dimethylformamide (DMF); *N,N*-dimethylacetamide (DMA); nitrobenzene; 1-methyl-2-pyrrolidinone. The addition of an appropriate base such as, for example, an alkali metal carbonate or hydrogen carbonate, sodium hydride or an organic base such as, for example, *N,N*-diethylethanamine or *N*-(1-methylethyl)-2-propanamine may be utilized to pick up the acid which is liberated during the course of the reaction. In some circumstances the addition of an iodide salt, preferably an alkali metal iodide, is appropriate. Somewhat elevated temperatures may enhance the rate of the reaction.

The compounds of formula (I) can also be prepared by the cyclodesulfurization reaction of an appropriate thiourea derivative of formula




Said cyclodesulfurization reaction may be carried out by the reaction of (IV) with an appropriate alkyl halide, preferably iodomethane in an appropriate reaction-inert organic solvent, e.g., a lower alkanol such as methanol, ethanol or 2-propanol. Otherwise, the cyclode-sulfurization reaction may be carried out by the



EP 0 151 824 B1

reaction of (IV) with an appropriate metal oxide or salt in an appropriate solvent according to art-known procedures. For example, the compounds of formula (I) can easily be prepared by the reaction of (IV) with an appropriate Hg(II) or Pb(II) oxide or salt, such as, for example HgO, HgCl₂, Hg(OAc)₂, PbO or Pb(OAc)₂. In certain instances it may be appropriate to supplement the reaction mixture with a small amount of sulfur. Even so methanediimines, especially *N,N*-methanetetraylbis[cyclohexanamine] may be used as cyclodesulfurizing agents.

The compounds of formula (I), wherein L is a radical of formula (b-1), wherein Y is NH and X is O or S, said X being represented by X¹, and wherein W is a radical of formula (c-1-c), said compounds being represented by the formula (I-b-1), can generally be prepared by reacting an isocyanate or isothiocyanate of formula (VI) with a reagent of formula (V):





The compounds of formula (I), wherein L is a radical of formula (b-1), wherein Y is other than a direct bond, said Y being Y¹, X is X¹, and wherein W is a radical of formula (c-1-c), wherein Z¹ is NH, said compounds being represented by the formula (I-b-2), or the compounds of formula (I), wherein L is a radical of formula (b-2), wherein X is X¹, and wherein W is a radical of formula (c-1-c), wherein Z¹ is NH, said compounds being represented by the formula (I-b-3), can be prepared by reacting an isocyanate or isothiocyanate of formula (VII) with an intermediate of formula (III-b-1), respectively (III-b-2).



The reaction of (V) with (VI), of (VII) with (III-b-1) or (III-b-2) is generally conducted in a suitable reaction-inert solvent, such as, for example, an ether, e.g., tetrahydrofuran. Elevated temperatures may be suitable to enhance the rate of the reaction. When W¹ is hydrogen, the reaction is conducted in aqueous medium containing an appropriate acid, such as, for example, acetic acid.


The compounds of formula (I) wherein L is a radical of formula (b-1), wherein Y is Y¹ and X is X¹ and wherein W is other than a radical of formula (c-1-c), said W being represented by W¹, and said compounds being represented by the formula (I-c-1), and the compounds of formula (I), wherein L is a radical of formula (b-2), wherein X is X¹ and W is W³, said compounds being represented by the formula (I-c-2), may be prepared by reacting an intermediate of formula (III-b-1) respectively (III-b-2) with as reagent of formula (VIII).



The reaction of (III-b-1) or (III-b-2) with (VIII) may generally be conducted following art-known esterification-


or amidation reaction procedures. For example, the carboxylic acid may be converted into a reactive derivative, e.g., an anhydride or a carboxylic acid halide, which subsequently, is reacted with (III-b-1) or (III-b-2); or by reacting (III-b-1) or (III-b-2) and (VIII) with a suitable reagent capable of forming amides or esters, e.g., dicyclohexylcarbodiimide, 2-chloro-1-methylpyridinium iodide. Said reactions are most conveniently conducted in a suitable solvent such as, for example, an ether, e.g. tetrahydrofuran, a halogenated hydrocarbon, e.g. dichloromethane, trichloromethane or a polar aprotic solvent, e.g. *N,N*-dimethylformamide. The addition of a base, e.g. *N,N*-diethylethanamine may be appropriate.

The compounds of formula (I) wherein L is a radical of formula (b-1), (b-3), (b-4), (b-5), (b-6) or (b-7), said compounds being represented by the formula (I-d), may also be prepared by reacting an appropriate alkenylene of formula (IX) with a piperidine of formula (III-a) by stirring and, if desired, heating the reactants together.




L¹ is selected so, that it forms, combined with —Alk—, a radical of formula (b-1), (b-3), (b-4), (b-5), (b-6) or (b-7).

The compounds of formula (I), wherein L is a radical of formula (b-5), wherein Alk is —CH₂—, wherein Het is a substituted 2-furanyl radical, said compounds being represented by the formula (I-e), can be prepared by reacting a substituted furan of formula (X) with an intermediate of formula (III-a) in the presence of formaldehyde or a polymeric form thereof in a suitable solvent.







wherein R¹⁸ is a previously described substituent of said furan ring.

The compounds of formula (I-e), wherein R¹⁸ is a radical of formula (c-4-d-1), wherein Y is Y¹, said compounds being represented by the formula (I-e-1), or the compounds of formula (I-e), wherein R¹⁸ is a radical of formula (c-4-d-1) wherein Z is Z¹, said compounds being represented by the formula (I-e-2), can be prepared by reacting an intermediate of formula (X-a), (X-b) (X-c) or (X-d) with a reagent of formula (XI-a), (XI-b), (XI-c) or (XI-d); in order to simplify the structural representation of the compounds of formula (I-e-1) and (I-e-2) and the intermediates of formula (X-a), (X-b), (X-c) and (X-d), the






radical will further be represented by the symbol D¹.







The compounds of formula (I), wherein L is a radical of formula (b-3), wherein Y^a is other than a direct bond, said Y^a being represented by Y^{a-1} , and said compounds by the formula (I-f-1), or wherein L is a radical of formula (b-4) wherein T is a radical of formula (c-3-a) or (c-3-b), said compounds being represented by the formula (I-f-2), respectively (I-f-3), can be prepared by reacting an intermediate of formula (XII-a), respectively (XII-b) and (XII-c), with a reagent of formula (XIII-a), respectively (XIII-b) and (XIII-c):



The reaction of the compounds of formulae (XI) with the compounds of formulae (X), and those of formulae (XIII) with those of formulae (XIII) is conveniently conducted following the same procedures as described hereinabove for the synthesis from (I) starting from (II) and (III).

The compounds of formula (I), wherein L is a radical of formula (b-3), wherein X^a is O or S, said X^a being X^{a-1} , and wherein Y^a is Y^{a-1} , and wherein W^2 is a radical of formula (c-2-a), wherein Z^1 is NH, said compounds being represented by the formula (I-g-1), and the compounds of formula (I), wherein L is a radical of formula (b-4), wherein T is a radical of formula (c-3-a), wherein X is Z^1 and Z is NH, said compounds being represented by the formula (I-g-2), can be prepared by reacting a reagent of formula (XIV-a) respectively (XIV-b) with an intermediate of formula (XII-a) respectively (XII-b).



The reaction of the compounds of formula (XIV) with those of formula (XII) can conveniently be conducted following the same procedures as described hereinabove for the reaction of (V) with (VI), and (VII) with (III-b-1) or (III-b-2).

The compounds of formula (I) can also be converted into each other following art-known procedures of functional group transformation. Some examples will be cited hereinafter.

The compounds of formula (I) having a nitro substituent can be converted into their corresponding


amines by stirring and, if desired, heating the starting nitro-compounds in a hydrogen-containing medium in the presence of a suitable amount of an appropriate catalyst such as, for example, platinum-on-charcoal, palladium-on-charcoal, Raney-nickel and the like catalysts. Suitable solvents are, for example, alcohols, e.g., methanol or ethanol.

5 Halo atoms substituted on aryl groups may be replaced by hydrogen following art-known hydrogenolysis procedures, i.e., by stirring and, if desired, heating the starting compounds in a suitable solvent under hydrogen atmosphere in the presence of an appropriate catalyst, e.g., palladium-on-charcoal and the like catalysts. Said halo atoms may also be replaced by a lower alkoxy or a lower alkylthio substituent by reacting the starting halo-compound with an appropriate alcohol or thioalcohol or, preferably, an alkali- or earth alkaline metal salt or an appropriate alcohol or thioalcohol in a suitable solvent. Said lower alkoxy or alkylthio substituents may be converted into alcohol or thiol groups by hydrolysing the starting lower alkoxy or alkylthio compounds in an acidic aqueous medium such as, for example, an aqueous hydrogen halide solution.

The compounds of formula (I), containing a Y, Y' or Y² group of formula NH can be converted into compounds of formula (I) wherein Y, Y' or Y² is NR¹⁶, R¹⁶ being other than hydrogen, by reacting the starting amine with an appropriate N-alkylating or N-acylating agent such as, for example, a lower alkyl or Ar²-lower alkyl halogenide, e.g. bromomethane, iodoethane, (chloromethyl)benzene and the like; or a carboxylic acid or a derivative thereof, e.g. an acid halide or an acid anhydride.

The compounds of formula (II), containing a Y, Y¹ or Y² group of formula NR¹⁶, wherein R¹⁶ is the previously described radical of formula —C(X)—R¹⁷, wherein X is O or S and R¹⁷ is lower alkylamino, or Ar¹lower alkylamino can be prepared by reacting the starting amine with an appropriate isocyanate or isothiocyanate.

The compounds of formula (I) wherein L is a radical of formula



may be converted into compounds of formula (I) wherein L is a radical of formula




by reacting the former compounds with an appropriate acid in the presence of a suitable solvent, e.g., water.

In all of the foregoing and in the following preparations, the reaction products may be isolated from the reaction mixture and, if necessary, further purified according to methodologies generally known in the art.


The compounds of formula (I) have basic properties and, consequently, they may be converted to their therapeutically active non-toxic acid addition salt forms by treatment with appropriate acids, such as, for example, inorganic acids, such as hydrohalic acid, e.g. hydrochloric, hydrobromic and the like, and sulfuric acid, nitric acid or phosphoric acid; or organic acids, such as, for example, acetic, propanoic, hydroxyacetic, 4-hydroxypropanoic, 2-oxopropanoic, ethanedioic, propanedioic, butanedioic, (Z)-2-butenedioic, (E)-2-butenedioic, 2-hydroxybutanedioic, 2,3-dihydroxybutanedioic, 2-hydroxy-1,2,3-propanetriolic, cyclyl-, methyl-, ethyl-, propyl-, butyl-, pentyl-, hexyl-, heptyl-, octyl-, nonyl-, decyl-, dodecyl-, tetradecyl-, hexadecyl-, octadecyl-, eicosyl-, and triacosyl-sulfonic acids, 4-methylbenzenesulfonic acid, cyclohexanesulfonic acid, 4-methylbenzenesulfonic acid, and 4-amino-2-hydroxybenzoic acid. Conversely the salt form can be converted by treatment with alkali into the free base form.

50 A number of intermediates and starting materials in the foregoing preparations are known compounds which may be prepared according to art-known methodologies of preparing said or similar compounds. A number of such preparation methods will be described hereinafter in more detail.

The intermediates of formula (III-a) can conveniently be prepared starting from a thiourea derivative of formula




wherein P is an appropriate protective group such as, for example, lower alkyloxycarbonyl, $\text{Ar}^2\text{-CH}_2\text{-O-CO-}$, $\text{Ar}^2\text{-CH}_2\text{-}$ and the like, by a cyclodesulfurization reaction following the same procedure as described hereinabove for the preparation of (I) starting from (IV) and, subsequently eliminating the protective group P in the thus obtained intermediate of formula



(XVI)


The elimination of the protective group P in (XVI) may generally be carried out following art-known procedures such as, for example, by hydrolysis in alkaline or acidic aqueous medium.

The intermediates of formula (III-b-1) and (III-c) may be derived from the corresponding intermediates of formula (III-a) by reacting the latter with a suitable reagent following art-known N-alkylating procedures. For example, intermediates of formula (III-b-1) wherein $\text{HY}^1\text{-Alk-}$ represents a radical of formula $\text{H}_2\text{N-CH}_2\text{-Alk'}$, (III-b-1-a), can also be prepared by reacting an intermediate of formula (III-a) with a nitrile of formula (XVII) following art-known N-alkylating procedures and subsequently converting the thus obtained nitrile (XVIII) into the corresponding amine (III-b-1-a) following art-known nitrile to amine reducing procedures, e.g., by catalytically hydrogenating procedures.




In (XVII), (XVIII) and (III-b-1-a) Alk' has the same meaning as Alk provided that one methylene function is missing.

The intermediates of formula (III-b-1) wherein $\text{HY}^1\text{-Alk-}$ represents a radical of formula $\text{HY}^1\text{-CH}_2\text{-CH}_2\text{-}$, (III-b-1-b), may also be prepared by the reaction of (III-a) with a reagent of formula (XIX) by stirring and, if desired, heating the reactants together in a suitable solvent.





The intermediates of formula (III-b-1) may be converted into an intermediate of formula (III-c) by converting the function HY^1 into an appropriate leaving group, e.g., where Y^1 is O, said intermediates being represented by the formula (III-b-1-c) by converting a hydroxy function into a chloro atom, with thionyl chloride or phosphoryl chloride.



The intermediates of formula (III-b-1-a) may also be derived from an appropriate corresponding carbonyl-oxidated form by reacting said carbonyl-oxidated form with hydroxylamine and reducing the thus obtained oxime following art-known methods, e.g., catalytic hydrogenation and the like reducing methods.

The intermediates of formula (III-b-1) or (III-b-2) may also be prepared by reacting a reagent containing both a protected Y^1 or NH function and a carbonyl function, by reacting said reagent with (III-a) and reducing the thus obtained intermediate following art-known procedures, e.g. catalytic hydrogenation, followed by an elimination reaction of the group protecting Y^1 . For example, the intermediates of formula (III-b-2), wherein D is substituted by a 4-piperidinyl radical, said compounds being represented by the formula (III-b-2-a), can be prepared by reacting a reagent of formula (XX) with (III-a) followed by an appropriate reduction, and subsequently eliminating the protective group P as described hereinabove:





The intermediates of formula (VI) can conveniently be prepared by converting the amino group in the compounds of formula (III-b-1-a) into an isocyanato or isothiocyanato group following art-known procedures, for example, by reacting said amino group with CS_2 in the presence of ethyl carbonochloridate.

The intermediates of formula (X-a) can be converted into intermediates of formula (X-b) by a suitable conversion of the Y^1H group into a leaving group; the intermediates of formula (X-c) wherein Y^1 is other than a direct bond, said Y^1 , can also be prepared by alkylating (X-a) with an appropriate reagent; the intermediates of formula (X-c) can be converted into those of formula (X-d) by a suitable conversion of the Z^1H group into a leaving group.

The intermediates of formula (XII-a), wherein Y^1 is other than a direct bond, can be prepared by alkylating an intermediate of formula (III-b-1) with an appropriate aromatic reagent; the intermediates of formula (XII-b) and (XII-c) can be prepared following art-known procedures as described in, for example, U.S. Patent No. 4,219,559.

The intermediates of formula (XV) and those of formula (XV) wherein R^2 is hydrogen, (XV-a), may be prepared by reacting a piperidine of formula (XXII-a) of (XXII-b) with an aromatic reagent of formula (XXIII-a) of (XXIII-b).



During one of the reactions the intermediates wherein R^1 and/or R^2 and/or R^8 and/or R^{15} and/or R^{16} is hydrogen may be converted into the corresponding intermediates wherein R^1 and/or R^2 and/or R^8 and/or R^{15} and/or R^{16} is other than hydrogen following art-known N-alkylating, N-acylating or reductive N-alkylating procedures.

From formula (I) it is evident that the compounds of this invention may have several asymmetric carbon atoms in their structure. Each of these chiral centers may be present in a R- and a S-configuration, this R- and S-notation being in correspondence with the rules described by J. Org. Chem. 35 (9), 2849—2867 (1970).

Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained by the application of art-known procedures. Diastereoisomers may be separated by physical separation methods such as selective crystallization and chromatographic techniques, e.g., counter current distribution, and enantiomers may be separated from each other by the selective crystallization of their diastereomeric salts with optically active acids.

Pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically.

It is evident that the cis and trans diastereomeric racemates may be further resolved into their optical isomers, cis-(+), cis-(−), trans-(+) and trans-(−) by the application of methodologies known to those skilled in the art.

Stereochemically isomeric forms of the compounds of formula (I) are naturally intended to be embraced within the scope of the invention.

The compounds of formula (I) have histamine antagonistic properties and some of the compounds of formula (I) have also serotonin-antagonistic properties.

EP 0 151 824 B1

The useful antihistaminic properties of the compounds of formula (I) are demonstrated in the following test procedure.

Protection of rats from compound 48/80-induced lethality.

5 Compound 48/80, a mixture of oligomers obtained by condensation of 4-methoxy-N-methylbenzeneethanamine and formaldehyde has been described as a potent histamine releasing agent (Int. Arch. Allergy, 13, 336 (1958)). The protection from compound 48/80-induced lethal circulatory collapse appears to be a simple way of evaluating quantitatively the antihistaminic activity of test compounds. Male
10 rats of an inbred Wistar strain, weighing 240—260 g were used in the experiment. After overnight starvation the rats were transferred to conditioned laboratories (temp. = $21 \pm 1^\circ\text{C}$, relative humidity = $65 \pm 5\%$). The rats were treated subcutaneously or orally with a test compound or with a solvent (NaCl solution, 0.9%). One hour after treatment there was injected intravenously compound 48/80, freshly dissolved in water, at a dose of 0.5 mg/kg (0.2 ml/100 g of body weight). In control experiments, wherein
15 250 solvent-treated animals were injected with the standard dose of compound 48/80, not more than 2.8% of the animals survived after 4 hours. Survival after 4 hours is therefore considered to be a safe criterion of a protective effect of drug administration.

The ED_{50} -values of the compounds of formula (I) are listed in table 1. Said ED_{50} -values are the values in mg/kg body weight at which the tested compounds protect 50% of the tested animals against compound
20 48/80-induced lethality.

The compounds listed in table 1 are given for the purpose of exemplifying the useful pharmacological activities of all the compounds within the scope of formula (I).

Table I (cont'd)


Comp. No.	L	R ¹	base or salt form	mp, °C	Column 1	
					Compound 48/80	lethality test in rate-ED ₅₀ in mg/kg body weight
					$-A^1-A^2-A^3-A^4-$	
25	$(CH_3)_2N-C-NH(CH_2)_2$ N-CN	4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- base	110.5	base	0.08
24	$CH_3O-CHCH_2NHCH_2(CH_2)_2$ N-CN	4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- (E)-2- butenedioate(1:2)	173.1	base	0.16
21	$CH_3S-C-NH(CH_2)_2$ N-CN	4-F-C ₆ H ₄ CH ₂	-N=CH-CH=CH- base	172.2	base	0.16
26	$O \begin{array}{c} \diagup \diagdown \\ \text{N} \end{array} (CH_2)_2 NHCH_2(CH_2)_2$ N-CN	4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- H ₂ O	125.6	base	0.16
35	$(C_2H_5O)-C-NH(CH_2)_2$ S	4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- base	148.6	base	0.16
34	$H_2N-NH-C-NH(CH_2)_2$ S	4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- H ₂ O	183.8	base	0.02
40		4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- base	162.7	base	0.16
28	$O \begin{array}{c} \diagup \diagdown \\ \text{N} \end{array} C-NH(CH_2)_2$ S	4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- base	191.6	base	0.16
31	$(CH_3)_2NH-C-NH(CH_2)_2$ S	4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- base	159.7	base	0.08

Table I (cont'd)

Comp. No.	L	R ¹	-A ¹ =A ² -A ³ =A ⁴ - base or salt form	mp. °C	Column 1	
					Compound 48/80	lethality test in rats-BD50 in mg/kg body weight
32		4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- base	175.5	0.16	
30		4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- H ₂ O	124.6	0.08	
36		4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- H ₂ O	152.4	0.16	
37		4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- base	218.8	0.08	
45		4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- base	153.2	0.08	
46		(2-furyl)CH ₂	-CH=CH-CH=CH- 1/2H ₂ O	125.2	0.16	
56		4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- base	186.9	0.04	
52		4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- (E)-2-butenedioate (1:2)	175.2	0.31	
51		4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- base	170.9	0.16	

* : Ac = acetyl

Table 1 (cont'd)













Comp. No.	L	R ¹	-A ¹ =A ² -A ³ =A ⁴ - base or salt form	mp. °C	Column 1	
					Compound 48/80	lethality test in rats-ED50 in mg/kg body weight
1	CH ₂ =CH-O-(CH ₂) ₂	4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- base	138.5		0.08
57	CH ₃ -NH-CO-NH-  -(CH ₂) ₂	4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- base	300		0.08
53	CH ₃ SO ₂ NH-  -(CH ₂) ₂	4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- H ₂ O	191.0		0.16
12	 -(CH ₂) ₂	4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH- base	168.4		0.16
14	 -(CH ₂) ₂	(2-furanyl)CH ₂	-CH=CH-CH=CH- H ₂ O	133.2		0.08
15	 -(CH ₂) ₂	(2-furanyl)CH ₂	-N=CH-CH=CH- base	171.5		0.01
16	 -(CH ₂) ₂	4-F-C ₆ H ₄ CH ₂	-N=CH-CH=CH- H ₂ O	167.1		0.04

Table 1 (cont'd)

Comp. No.	L	R ¹	1 2 3 4 -A =A =A =A -	base or salt form	mp. °C	Column 1
						Compound 48/80 lethality test in rats-ED50 in mg/kg body weight
20		4-F-C ₆ H ₄ CH ₂	-CH=CH-CH=CH-	base	170.5	0.31
6		(4-thiazolyl)CH ₂	-CH=CH-CH=CH-	2HCl 1 1/2 H ₂ O	187.2	0.16
7		(2-pyridinyl)CH ₂	-N=CH-CH=CH-	3HCl 2 H ₂ O	190.6	0.04
8		(2-thienyl)CH ₂	-CH=CH-CH=CH-	base	167.6	0.16
9		(2-pyridinyl)CH ₂	-CH=CH-CH=CH-	2HCl 2 H ₂ O	185.1	0.08
10		(3-pyridinyl)CH ₂	-CH=CH-CH=CH-	H ₂ O	147.3	0.31

In view of their antihistaminic properties, the compounds of formula (I) and their acid-addition salts are very useful in the treatment of allergic diseases such as, for examples, allergic rhinitis, allergic conjunctivitis, chronic urticaria or allergic asthma.

In view of their useful antihistaminic properties, the subject compounds may be formulated into various pharmaceutical forms for administration purposes.

To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, in base or acid-addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration.

These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for administration orally, rectally or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils and alcohols in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders or disintegrating agents in the case of powders, pills, capsules and tablets.

Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers or suspending agents may be employed.

Acid addition salts of (I) due to their increased water solubility over the corresponding base form, are obviously more suitable in the preparation of aqueous compositions.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form of ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls or tablespoonfuls, and segregated multiples thereof.

The present invention is also related with a method of preparing a medicament for treating allergic diseases in warm-blooded animals suffering from said allergic diseases suitable for administering an effective anti-allergic amount of a compound of formula (I) or a pharmaceutically acceptable acid addition salt thereof.

Suitable doses administered daily to subjects are varying from 0.1 to 110 mg, more preferably from 1 to 50 mg.

The following examples are intended to illustrate the scope of the present invention. Unless otherwise stated all parts therein are by weight.

EXAMPLES

A. Preparation of Intermediates:

Example 1

A mixture of 90 parts of 4-chloro-3-nitropyridine, 71 parts of 4-fluorobenzenemethanamine, 63 parts of sodium carbonate and 900 parts of *N,N*-dimethylacetamide was stirred for 1 hour at 50°C. Water was added and the product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was crystallized from acetonitrile. The product was filtered off and dried, yielding 106 parts (75%) of *N*-[(4-fluorophenyl)methyl]-3-nitro-4-pyridinamine; mp. 136.8°C (intermediate 1).

In a similar manner there were also prepared:

N-[(4-fluorophenyl)methyl]-2,3-pyridinediamine as a residue (2);
N-[(4-fluorophenyl)methyl]-4-nitro-3-pyridinamine, 1-oxide (3); 2-nitro-*N*-(2-thienylmethyl)-benzenamine (4);
N-(3-nitro-2-pyridinyl)-2-pyridinemethanamine; mp. 113.6°C (5); and 3-nitro-*N*-(2-thienylmethyl)-2-pyridinamine; mp. 100°C (6).

Example 2

To a stirred and cooled (0°C) solution of 8.7 parts of *N*-[(4-fluorophenyl)methyl]-4-nitro-3-pyridinamine, 1-oxide and 150 parts of trichloromethane was added dropwise a solution of 10.2 parts of phosphorus trichloride in 75 parts of trichloromethane. Upon completion, the mixture was allowed to reach room temperature and stirring was continued for 1 hour at reflux temperature. The reaction mixture was cooled and the solvent was evaporated. The residue was stirred in trichloromethane. The product was filtered off and dried, yielding 9 parts of *N*-[(4-fluorophenyl)methyl]-4-nitro-3-pyridinamine monohydrochloride (7).

Example 3

A mixture of 11 parts of *N*-[(4-fluorophenyl)methyl]-4-nitro-3-pyridinamine monohydrochloride, 2 parts

of a solution of thiophene in ethanol 4% and 240 parts of methanol saturated with ammonia was hydrogenated at normal pressure and at room temperature with 3 parts of platinum-on-charcoal catalyst 5%. After the calculated amount of hydrogen was taken up, the whole was warmed and the catalyst was filtered off. It was washed with 2-methoxyethanol. The filtrate was evaporated and the residue was heated in acetonitrile. After stirring and cooling, the product was filtered off and dried, yielding 6.5 parts (58%) of N⁴[[4-(4-fluorophenyl)methyl]-3,4-pyridinediamine monohydrochloride; mp. 208.9°C (8).

In a similar manner there were also prepared:

N²-(2-furanylmethyl)-2,3-pyridinediamine as a residue (9);

N⁴[[4-(4-fluorophenyl)methyl]-3,4-pyridinediamine; mp. 163.7°C (10).

N¹-(2-thienylmethyl)-1,2-benzenediamine (11);

N²-(2-pyridinylmethyl)-2,3-pyridinediamine; mp. 134.9°C (12);

N²-(2-thienylmethyl)-2,3-pyridinediamine; mp. 92.1°C (13).

Example 4

To a stirred and cooled mixture of 4 parts of sodium hydroxide in 60 parts of water were added successively 7.9 parts of carbon disulfide and 17.2 parts of ethyl 4-amino-1-piperidinecarboxylate at a temperature below 10°C. Stirring was continued for 30 minutes at this temperature. Then there were added dropwise 10.9 parts of ethyl carbonochloridate (exothermic reaction: temp. rises to about 35°C). Upon completion, stirring was continued for 2 hours at 60°C. The reaction mixture was cooled and the product was extracted with methylbenzene. The extract was dried, filtered and evaporated, yielding 22 parts (100%) of ethyl 4-isothiocyanato-1-piperidine-carboxylate as a residue (14).

Example 5

A mixture of 84.7 parts of ethyl 4-isothiocyanato-1-piperidinecarboxylate, 86.8 parts of N⁴[[4-(4-fluorophenyl)methyl]-3,4-pyridinediamine and 450 parts of tetrahydrofuran was stirred and refluxed for 3 hours. The reaction mixture was evaporated and the residue was crystallized from acetonitrile, yielding 90 parts (52%) of ethyl 4-[[[4-[[4-(4-fluorophenyl)methyl]amino]-3-pyridinyl]-amino]thioxomethyl]amino]-1-piperidinecarboxylate; mp. 166°C (15).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there were also prepared:

ethyl 4-[[[2-[(2-furanylmethyl)amino]phenyl]amino]thioxomethyl]-amino]-1-piperidinecarboxylate as a residue (16);

ethyl 4-[[[3-[[4-(4-fluorophenyl)methyl]amino]-2-pyridinyl]amino]thioxomethyl]amino]-1-piperidinecarboxylate as a residue (17);

ethyl 4-[[[2-[(2-furanylmethyl)amino]-3-pyridinyl]aminothioxomethyl]amino]-1-piperidinecarboxylate; mp. 132.7°C (18);

ethyl 4-[[[3-[[4-(4-fluorophenyl)methyl]amino]-4-pyridinyl]aminothioxomethyl]amino]-1-piperidinecarboxylate as a residue (19).

ethyl 4-[[[2-[(2-thienylmethyl)amino]phenyl]aminothioxomethyl]amino]-1-piperidinecarboxylate as a residue (20).

ethyl 4-[[[2-[(2-pyridinylmethyl)amino]-3-pyridinyl]aminothioxomethyl]amino]-1-piperidinecarboxylate as a residue (21).

ethyl 4-[[[2-[(2-thienylmethyl)amino]-3-pyridinyl]aminothioxomethyl]amino]-1-piperidinecarboxylate as a residue (22).

Example 6

A mixture of 74 parts of ethyl 4-[[[2-[(2-furanylmethyl)amino]-3-pyridinyl]aminothioxomethyl]amino]-1-piperidinecarboxylate, 96 parts of mercury (II) oxide, 0.1 parts of sulfur and 800 parts of ethanol was stirred and refluxed for 3 hours. The reaction mixture was filtered over Hyflo and the filtrate was evaporated. The residue was crystallized from acetonitrile, yielding 52.5 parts (79%) of ethyl 4-[[[3-[(2-furanylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]amino]-1-piperidinecarboxylate; mp. 149.2°C (23).

In a similar manner there were also prepared:

ethyl 4-[[[1-[(2-furanylmethyl)-1H-benzimidazol-2-yl]amino]-1-piperidinecarboxylate; mp. 135.8°C (24);

ethyl 4-[[[1-[(4-fluorophenyl)methyl]-1H-imidazo[4,5-b]pyridin-2-yl]amino]-1-piperidinecarboxylate; mp. 212.5°C (25);

ethyl 4-[[[1-[(4-fluorophenyl)methyl]-1H-imidazo[4,5-c]pyridin-2-yl]amino]-1-piperidinecarboxylate dihydrochloride monohydrate (26);

ethyl 4-[[[3-[(4-fluorophenyl)methyl]-3H-imidazo[4,5-c]pyridin-2-yl]-amino]-1-piperidinecarboxylate dihydrochloride monohydrate; mp. 168.6°C (27);

ethyl 4-[[[1-[(2-thienylmethyl)-1H-benzimidazol-2-yl]amino]-1-piperidinecarboxylate; mp. 142.7°C (28);

ethyl 4-[[[3-[(2-pyridinylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]amino]-1-piperidinecarboxylate; mp. 141.3°C (29); and

ethyl 4-[[[3-[(2-thienylmethyl)-3H-imidazo[4,5-b]pyridin-2-yl]amino]-1-piperidinecarboxylate as a residue (30).

Example 7

A mixture of 14.5 parts of ethyl 4-[(1*H*-benzimidazol-2-ylamino)-1-piperidinecarboxylate, 13 parts of 2-(chloromethyl)pyrazine, 10.5 parts of sodium carbonate and 135 parts of *N,N*-dimethylformamide was stirred and heated for 3 hours at 50°C. The whole was further stirred overnight at 70°C. The reaction mixture was cooled and poured onto water. The product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the hydrobromide salt in 2-propanone. The salt was filtered off and dried, yielding 8.7 parts (32%) of ethyl 4-[[1-(2-pyrazinylmethyl)-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate dihydrobromide monohydrate; mp. 178.5—179.3°C (31).

In a similar manner there were also prepared:

ethyl 4-[[1-(4-thiazolylmethyl)-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate; mp. 156.2°C (32);
ethyl 4-[[1-(3-pyridinylmethyl)-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate; mp. 191.4°C (33);

and
ethyl 4-[[1-(2-pyridinyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate; mp. 161.5°C (34).

Example 8

A mixture of 50 parts of ethyl 4-[[3-(2-furanyl)methyl]-3*H*-imidazo[4,5-*b*]pyridin-2-yl]amino]-1-piperidinecarboxylate, 50 parts of potassium hydroxide, 400 parts of 2-propanol and 20 drops of water was stirred and refluxed for about 5 hours. The reaction mixture was evaporated and water was added to the residue. The product was extracted twice with 4-methyl-2-pentanone. The combined extracts were dried, filtered and evaporated. The solid residue was stirred in 1,1'-oxybisethane. The product was filtered off and dried, yielding 34 parts (85%) of 3-(2-furanyl)methyl)-*N*-(4-piperidinyl)-3*H*-imidazo[4,5-*b*]pyridin-2-amine; mp. 159.0°C (35).

In the similar manner there were also prepared:

1-(2-furanyl)methyl)-*N*-(4-piperidinyl)-1*H*-benzimidazol-2-amine; mp. 211.0°C (36);
N-(4-piperidinyl)-1-(2-thienylmethyl)-1*H*-benzimidazol-2-amine (37); and
N-(4-piperidinyl)-3-(2-thienylmethyl)-3*H*-imidazo[4,5-*b*]pyridin-2-amine; mp. 189.6—193.5°C (38).

Example 9

A mixture of 23.8 parts of ethyl 4-[[3-[(4-fluorophenyl)methyl]-3*H*-imidazo[4,5-*c*]pyridin-2-yl]amino]-1-piperidine carboxylate and 275 parts of a hydrobromic acid solution 48% in water was stirred overnight at 80°C. The reaction mixture was evaporated and the residue was crystallized from ethanol, yielding 14.7 parts (48%) of 3-[(4-fluorophenyl)methyl]-*N*-(4-piperidinyl)-3*H*-imidazo[4,5-*c*]pyridin-2-amine dihydrobromide monohydrate; mp. 291.6°C (39).

In a similar manner there were also prepared:

1-[(4-fluorophenyl)methyl]-*N*-(4-piperidinyl)-1*H*-imidazo[4,5-*b*]pyridin-2-amine dihydrobromide; mp. + 300.6°C (40);
1-[(4-fluorophenyl)methyl]-*N*-(4-piperidinyl)-1*H*-imidazo[4,5-*c*]pyridin-2-amine dihydrobromide; mp. + 279.4°C (41);
N-(4-piperidinyl)-1-(4-thiazolylmethyl)-1*H*-benzimidazol-2-amine dihydrobromide monohydrate; mp. 223.5°C (42);
N-(4-piperidinyl)-1-(2-pyrazinylmethyl)-1*H*-benzimidazol-2-amine trihydrobromide; (43);
N-(4-piperidinyl)-1-(3-pyridinylmethyl)-1*H*-benzimidazol-2-amine trihydrobromide; mp. >260°C (44);
N-(4-piperidinyl)-3-(2-pyridinylmethyl)-3*H*-imidazo[4,5-*b*]pyridin-2-amine trihydrobromide; mp. 265.5°C (45); and
N-(4-piperidinyl)-1-[(2-pyridinyl)methyl]-1*H*-benzimidazol-2-amine trihydrobromide; mp. 295.9°C (46);

Example 10

A mixture of 8.62 parts of 2-chloroacetonitrile, 37 parts of (*cis*+*trans*)-1-[(4-fluorophenyl)methyl]-*N*-(3-methyl-4-piperidinyl)-1*H*-benzimidazol-2-amine, 15.9 parts of sodium carbonate and 270 parts of *N,N*-dimethylformamide was stirred for 2 hours at 40°C. The reaction mixture was poured onto water. The product was extracted twice with 4-methyl-2-pentanone. The combined organic layers were dried, filtered and evaporated. The residue was crystallized from 1,1'-oxybisethane. The product was filtered off and dried, yielding 25.1 parts (57%) of (*cis*+*trans*)-4-[[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-3-methyl-1-piperidineacetonitrile; mp. 150.1°C (47).

In a similar manner there were also prepared:

4-[[1-(2-furanyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidine-acetonitrile; mp. 194.4°C (48); and
4-[[3-[(4-fluorophenyl)methyl]-3*H*-imidazo[4,5-*b*]pyridin-2-yl]amino]-1-piperidineacetonitrile; mp. 183.7°C (49).

Example 11

A mixture of 15 parts of 4-[[3-[(4-fluorophenyl)methyl]-3*H*-imidazo[4,5-*b*]pyridin-2-yl]amino]-1-piperidineacetonitrile and 400 parts of methanol saturated with ammonia was hydrogenated at normal

pressure and at room temperature with 3 parts of Raney-nickel catalyst. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated. The residue was crystallized from a mixture of acetonitrile and 2,2'-oxybispropane, yielding 10 parts (68%) of *N*-[1-(2-aminoethyl)-4-piperidinyl]-3-[(4-fluorophenyl)methyl]-3*H*-imidazo[4,5-*b*]pyridin-2-amine; mp. 174.5°C (50).

In a similar manner there were also prepared:

N-[1-(2-aminoethyl)-4-piperidinyl]-1-(2-furanyl)methyl]-1*H*-benzimidazol-2-amine; mp. 163.0°C (51);
cis-trans-*N*-[1-(2-aminoethyl)-3-methyl-4-piperidinyl]-1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-amine; mp. 132.2°C (52).

Example 12

A mixture of 9 parts of oxirane, 3.24 parts of 1-(4-fluorophenyl)methyl)-*N*-(4-piperidinyl)-1*H*-benzimidazol-2-amine and 400 parts of methanol was stirred first overnight at room temperature and further for 4 hours at 50°C. The reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated.

The residue was crystallized from a mixture of 4-methyl-2-pentanone and 2,2'-oxybispropane, yielding 15 parts of 4-[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-ylamino]-1-piperidineethanol; mp. 138.7°C (53).

Example 13

To 2 parts of a solution of 2 parts of thiophene in 40 parts of ethanol were added 15 parts of ethyl 4-oxo-1-piperidinecarboxylate, 25 parts of 1-(4-fluorophenyl)methyl)-*N*-(4-piperidinyl)-1*H*-benzimidazol-2-amine, and 200 parts of methanol. The whole was hydrogenated at normal pressure and at room temperature with 5 parts of platinum-on-charcoal catalyst 5%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the hydrochloride salt in 2-propanol and 2-propanone. The salt was filtered off and dried, yielding 13.6 parts of ethyl 4-[1-(4-fluorophenyl)methyl)-1*H*-benzimidazol-2-ylamino] [1,4'-bipiperidine]-1'-carboxylate dihydrochloride monohydrate; mp. 260°C (54).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there was also prepared: 1-[(4-fluorophenyl)methyl)-*N*-[1'-(phenylmethyl)-[1,3'-bipiperidin]-4-yl]-1*H*-benzimidazol-2-amine; mp. 174.6°C (55).

Example 14

A mixture of 21 parts of ethyl 4-[1-(4-fluorophenyl)methyl)-1*H*-benzimidazol-2-ylamino] [1,4'-bipiperidine]-1'-carboxylate and 450 parts of hydrobromic acid solution 48% was stirred and refluxed for 16 hours. The reaction mixture was evaporated. From the residue the free base was liberated in the conventional manner with sodium hydroxide in water and extracted with dichloromethane. The extract was dried, filtered and evaporated, yielding 8 parts (50%) of *N*-[1-(4-fluorophenyl)methyl)-1*H*-benzimidazol-2-yl]-[1,4'-bipiperidine]-4-amine as a residue (56).

Example 15

A mixture of 11.3 parts of 1-[(4-fluorophenyl)methyl)-*N*-[1'-(phenylmethyl)-[1,3'-bipiperidin]-4-yl]-1*H*-benzimidazol-2-amine and 200 parts of methanol was hydrogenated at normal pressure and at room temperature with 2 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated. The residue was suspended in 2,2'-oxybispropane. The product was filtered off and dried, yielding 8.5 parts (91.5%) of *N*-[1-(3'-bipiperidin)-4-yl]-1-[(4-fluorophenyl)methyl)-1*H*-benzimidazol-2-amine (57).

Example 16

To a stirred and hot (50°C) mixture of 4.1 parts of 2*H*-3,1-benzoxazine-2,4(1*H*)-dione and 31.5 parts of *N,N*-dimethylformamide was added dropwise a solution of 9.4 parts of *N*-[1-(2-aminoethyl)-4-piperidinyl]-1-[(4-fluorophenyl)methyl)-1*H*-benzimidazol-2-amine in 31.5 parts of *N,N*-dimethylformamide at 50°C. Upon completion, stirring was continued for 3 hours at 50°C. Water was added and the product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was crystallized from acetonitrile, yielding 9.8 parts (80%) of 2-amino-*N*-[2-[4-[1-(4-fluorophenyl)-methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]benzamide; mp. 171.7°C (58).

In a similar manner there was also prepared:
 2-(ethylamino)-*N*-[2-[4-[1-(4-fluorophenyl)methyl)-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]benzenamide monohydrate; mp. 139.8°C (59).

Example 17

To a stirred solution of 3 parts of 3-(2-hydroxyethyl)-2,4-(1*H*,3*H*)-pyrimidinedione and 45 parts of trichloromethane were added dropwise 8 parts of thionyl chloride. Upon completion, stirring was continued for 1 hour at reflux temperature. The reaction mixture was cooled. The precipitated product was filtered off and crystallized from 2-propanol, yielding 3.1 parts of 3-(2-chloroethyl)-2,4-(1*H*,3*H*)-pyrimidinedione; mp. 170°C (60).

B. Preparation of Final Compounds:

Example 18

A mixture of 1.6 parts of 1-chloro-2-(ethenylloxy)ethane, 7.3 parts of 1-(4-fluorophenyl)methyl)-*N*-(4-piperidinyl)-1*H*-benzimidazol-2-amine dihydrobromide, 3.1 parts of sodium carbonate, 0.1 parts of potassium iodide and 135 parts of *N,N*-dimethylformamide was stirred and heated overnight at 70°C. The reaction mixture was poured onto water and the product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile, yielding 1.9 parts (32%) of *N*-[1-[2-(ethenylloxy)ethyl]-4-piperidinyl]-1-(4-fluorophenyl)methyl)-1*H*-benzimidazol-2-amine; mp. 138.5°C (compound 1).

In a similar manner there were also prepared:

- 1-[1-(4-fluorophenyl)methyl)-*N*-[1-[2-(4-methoxyphenyl)]ethyl-4-piperidinyl]-1*H*-imidazo[4,5-*b*]pyridin-2-amine; mp. 186.8°C (compound 2);
- 1-[1-(4-fluorophenyl)methyl)-*N*-[1-[2-(4-methoxyphenyl)]ethyl-4-piperidinyl]-1*H*-imidazo[4,5-*c*]pyridin-2-amine; mp. 184.5°C (compound 3);
- 3-[1-(4-fluorophenyl)methyl)-*N*-[1-[2-(4-methoxyphenyl)]ethyl-4-piperidinyl]-3*H*-imidazo[4,5-*c*]pyridin-2-amine (E)-2-butenedioate (1:2); mp. 202.8°C (compound 4);
- 3-[2-[4-[[1-[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-2,4-(1*H*,3*H*)-pyrimidinedione; mp. 245.8°C (compound 5);
- 3-[2-[4-[[1-[1-(4-thiazolyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-2*H*-1-benzopyran-2-one dihydrochloride.sesquihydrate; mp. 187.2°C (compound 8);
- 3-[2-[4-[[3-[2-pyridinylmethyl]-3*H*-imidazo[4,5-*b*]pyridin-2-yl]amino]-1-piperidinyl]ethyl]-2*H*-1-benzopyran-2-one trihydrochloride.dihydrate; mp. 190.6°C (compound 7);
- 3-[2-[4-[[1-[2-thienylmethyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-2*H*-1-benzopyran-2-one; mp. 167.6°C (compound 8);
- 3-[2-[4-[[1-[2-pyridinylmethyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-2*H*-1-benzopyran-2-one dihydrochloride.dihydrate; mp. 185.1°C (compound 9);
- 3-[2-[4-[[1-[3-pyridinylmethyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-2*H*-1-benzopyran-2-one monohydrate; mp. 147.3°C (compound 10);
- 3-[2-[4-[[1-[2-thienylmethyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-2*H*-1-benzopyran-2-one; mp. 164.6°C (compound 11)

Example 19

A mixture of 3.8 parts of 3-(2-bromoethyl)-2*H*-1-benzopyran-2-one, 7.3 parts of 1-(4-fluorophenyl)methyl)-*N*-(4-piperidinyl)-1*H*-benzimidazol-2-amine dihydrobromide, 4.8 parts of sodium carbonate and 135 parts of *N,N*-dimethylformamide was stirred and heated overnight at 70°C. The reaction mixture was poured onto water. The product was extracted with trichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile, yielding 1.6 parts (21.5%) of 3-[2-[4-[[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-2*H*-1-benzopyran-2-one; mp. 168.4°C (compound 12).

In a similar manner there was also prepared:

- N*-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]-1-(2-pyrazinyl)methyl)-1*H*-benzimidazol-2-amine; mp. 127.4°C (compound 13);
- 3-[2-[4-[[1-(2-furanyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-2*H*-1-benzopyran-2-one monohydrate; mp. 133.2°C (compound 14);
- 3-[2-[4-[[3-(2-furanyl)methyl]-3*H*-imidazo[4,5-*b*]pyridin-2-yl]amino]-1-piperidinyl]ethyl]-2*H*-1-benzopyran-2-one; mp. 171.5°C (compound 15); and
- 3-[2-[4-[[3-[[1-(4-fluorophenyl)methyl]-3*H*-imidazo[4,5-*b*]pyridin-2-yl]amino]-1-piperidinyl]ethyl]-2*H*-1-benzopyran-2-one monohydrate; mp. 167.1°C (compound 16).

Example 20

A mixture of 4.7 parts of 1-(2-chloroethyl)-4-methoxybenzene, 14 parts of *N*-(4-piperidinyl)-1-(4-thiazolyl)methyl)-1*H*-benzimidazol-2-amine dihydrobromide.monohydrate, 15 parts of sodium carbonate, 0.3 parts of sodium iodide and 90 parts of *N,N*-dimethylacetamide was stirred overnight at 80°C. Water was added and the product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The oily residue was converted into the hydrobromide salt in ethanol. The salt was filtered off and dried, yielding 9 parts of *N*-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]-1-(4-thiazolyl)methyl)-1*H*-benzimidazol-2-amine dihydrobromide. dihydrate; mp. 239.2°C (compound 17).

Example 21

To a stirred mixture of 4 parts of *N*-[1-(4-fluorophenyl)methyl)-1*H*-benzimidazol-2-yl]-[1,4'-bipiperidine]-

4-amine, 1 part of *N,N*-diethylethanamine and 91 parts of dichloromethane was added dropwise a solution of 1.6 parts of 4-fluorobenzoyl chloride in 39 parts of dichloromethane: slightly exothermic reaction, the temperature rises from 25°C to 30°C. Upon completion, stirring was continued for one hour at room temperature. The reaction mixture was purified by high performance liquid chromatography using a mixture of trichloromethane, hexane and methanol (45:45:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated, yielding 1.8 parts (34%) of 1'-[4-fluorobenzoyl-*N*-(1-(4-fluorophenyl)methyl)-1*H*-benzimidazol-2-yl)]-[1,4'-bipiperidine]-4-amine; mp. 194.3°C; (compound 18).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there was also prepared:

N,N-diethyl-4-[1-(4-fluorophenyl)methyl-1*H*-benzimidazol-2-ylamino]-[1,4'-bipiperidine]-1'-carboxamide; mp. 176.6°C (compound 19).

Example 22

A mixture of 1.64 parts of 2-methyl-1*H*-imidazole, 9.2 parts of *N*-[1-(2-chloroethyl)-4-piperidinyl]-1-(4-fluorophenyl)methyl-1*H*-benzimidazol-2-amine dihydrochloride, 6.4 parts of sodium carbonate and 135 parts of *N,N*-dimethylformamide was stirred overnight at 60°C. The reaction mixture was poured into water. The product was extracted with trichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile. The product was filtered off and dried, yielding 2.6 parts (30%) of 1-[(4-fluorophenyl)methyl-*N*-[1-(2-(2-methyl-1*H*-imidazol-1-yl)ethyl]-4-piperidinyl]-1*H*-benzimidazol-2-amine; mp. 170.5°C (compound 20).

Example 23

A mixture of 1.9 parts of dimethyl cyanocarbonimidodithioate, 4.8 parts of *N*-[1-(2-aminoethyl)-4-piperidinyl]-3-[(4-fluorophenyl)methyl-3*H*-imidazo[4,5-*b*]pyridin-2-amine and 80 parts of methanol was stirred for 2 hours at room temperature. The reaction mixture was evaporated and the residue was crystallized from acetonitrile, yielding 4.5 parts (74%) of *S*-methyl *N'*-cyano-*N*-[2-[4-[(3-[(4-fluorophenyl)methyl-3*H*-imidazo[4,5-*b*]pyridin-2-yl]amino]-1-piperidinyl)ethyl]carbamimidodithioate; mp. 172.2°C (compound 21).

In a similar manner there was also prepared:

S-methyl *N'*-cyano-*N*-[2-[4-[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl)ethyl]carbamimidodithioate (compound 22).

Example 24

A mixture of 1.5 parts of *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine, 3.7 parts of *N*-[1-(2-aminoethyl)-4-piperidinyl]-1-(4-fluorophenyl)methyl-1*H*-benzimidazol-2-amine and 80 parts of ethanol 50% was stirred overnight at room temperature. The reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from 2-propanol. The product was filtered off and dried overnight at 110°C, yielding 1.5 parts (33%) of *N*-[2-[4-[1-[(4-fluorophenyl)methyl-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl)ethyl]-*N'*-nitroguanidine; mp. 146.7°C (compound 23).

Example 25

A mixture of 1.6 parts of 2,2-diethoxyethanamine, 4.6 parts of *S*-methyl *N'*-cyano-*N*-[2-[4-[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl)ethyl]-carbamimidodithioate and 40 parts of 1-butanol was stirred and refluxed overnight. The reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (93:7 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (E)-2-butenedioate salt in ethanol. The salt was filtered off and dried, yielding 2 parts of *N'*-cyano-*N*-(2,2-dimethoxyethyl)-*N'*-[2-[4-[1-[(4-fluorophenyl)methyl-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl)ethyl]guanidine (E)-2-butenedioate (1:2) (compound 24).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there were also prepared:

N'-cyano-*N*-[2-[4-[1-[(4-fluorophenyl)methyl-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl)ethyl]-*N'*-dimethylguanidine; mp. 110.5°C (compound 25).

N'-cyano-*N*-[2-[4-[1-[(4-fluorophenyl)methyl-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl)ethyl]-*N'*-[2-(4-morpholinyl)ethyl]guanidine monohydrate; mp. 125.6°C (compound 26).

Example 26

A solution of 5.71 parts of (cis+trans)-*N*-[1-(2-aminoethyl)-3-methyl-4-piperidinyl]-1-(4-fluorophenyl)methyl-1*H*-benzimidazol-2-amine, 2.84 parts of 1,1'-thiocarbonylbis[1*H*-imidazole] in 180 parts of tetrahydrofuran was stirred for 2 hours at room temperature. 0.9 Parts of gaseous *N*-methylmethanamine was bubbled, during 30 minutes, through the mixture. Stirring was continued overnight at room temperature.

The whole was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (from 100:0 to 90:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was stirred in 2,2'-oxybispropane. The product was filtered off and dried, yielding 2.3 parts (32.7%) of *cis-N*-[2-[4-[[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-3-methyl-1-piperidinyl]ethyl]-*N,N'*-dimethylthiourea; mp. 126.7°C (compound 27).

In a similar manner there was also prepared:

N-[2-[4-[[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-4-morpholinecarbothioamide; mp. 191.6°C (compound 28).

Example 27

A mixture of 0.9 parts of piperidine, 4.1 parts of 1-(4-fluorophenyl)methyl-*N*-[1-(2-isothiocyanatoethyl)-4-piperidinyl]-1*H*-benzimidazol-2-amine and 135 parts of tetrahydrofuran was stirred for 2 hours at room temperature. The reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile, yielding 1 part (20.2%) of *N*-[2-[4-[[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-1-piperidine-carbothioamide; mp. 175.6°C (compound 29).

Example 28

A mixture of 3.75 parts of 3-amino-1-propanol, 20.5 parts of 1-(4-fluorophenyl)methyl-*N*-[1-(2-isothiocyanatoethyl)-4-piperidinyl]-1*H*-benzimidazol-2-amine and 450 parts of tetrahydrofuran was stirred for 3 hours at room temperature. The reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile, yielding 16 parts (64%) of *N*-[2-[4-[[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-*N'*-(3-hydroxypropyl) thiourea monohydrate; mp. 124.6°C (compound 30).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there were also prepared:

N-[2-[4-[[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-*N,N'*-dimethylthiourea; mp. 159.7°C (compound 31).

N,N-diethyl-*N'*-(2-[4-[[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl)thiourea; mp. 175.5°C (compound 32).

N'-(2-[4-[[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl)-*N*-(2-phenylethyl)thiourea (E)-2-butenedioate (1:2); mp. 196.8°C (compound 33).

N'-(2-[4-[[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl)hydrazinecarbothioamide monohydrate; mp. 183.8°C (compound 34).

Example 29

A mixture of 1.3 parts of 2-chloro-3-pyridinamine, 4.1 parts of 1-(4-fluorophenyl)methyl-*N*-[1-(2-isothiocyanatoethyl)-4-piperidinyl]-1*H*-benzimidazol-2-amine and 80 parts of ethanol was stirred and refluxed overnight. The reaction mixture was evaporated. Water and ammonia were added to the residue and the product was extracted with trichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile. The product was filtered off and dried, yielding 1.4 parts of ethyl [2-[4-[[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]carbamothioate; mp. 148.6°C (compound 35).

Example 30

A mixture of 0.55 parts of isocyanatomethane, 4 parts of *N*-[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]-[1,4'-bipiperidine]-4-amine, 80 parts of ethanol and 65 parts of dichloromethane was stirred for 3 hours at room temperature. The reaction mixture was evaporated. The residue was purified by HPLC over silica gel using a mixture of trichloromethane, hexane and methanol, saturated with ammonia, (45:45:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile, yielding 1 part (25%) of 4-[[1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-ylamino]-*N*-methyl-1,4'-bipiperidine)-1'-carboxamide monohydrate; mp. 152.4°C (compound 36).

Example 31

A mixture of 0.8 parts of isothiocyanatomethane, 4 parts of *N*-[1,3'-bipiperidin-4-yl]-1-[[4-fluorophenyl)methyl]-1*H*-benzimidazol-2-amine and 90 parts of tetrahydrofuran was stirred overnight at room temperature. The reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile, yielding 3.7 parts (77%) of 4-[[1-(4-fluorophenyl)methyl]-1*H*-

benzimidazol-2-yl]amino)-*N*-methyl-1,3'-bipiperidine]-1'-carbothioamide; mp. 218.8°C (compound 37).

In a similar manner there were also prepared:

4-[[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino)-*N*-methyl-1,4'-bipiperidine]-1'-carboxamide; mp. 222.7°C (compound 38).

5 *N*-cyclohexyl-*N'*-[2-4-[[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethylthiourea; mp. 177°C (compound 39).

N-[2-4-[[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl-*N'*phenylthiourea; mp. 162.7°C (compound 40).

10 *N*-[2-4-[[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-ylamino]-1-piperidinyl]ethyl]-*N'*-(4-methoxyphenyl)thiourea; mp. 165.9°C (compound 41).

Example 32

To a stirred mixture of 1.9 parts of 2-oxo-2*H*-benzopyran-3-carboxylic acid, 4.04 parts of *N,N*-diethylethanamine and 195 parts of dichloromethane were added 2.55 parts of 2-chloro-1-methylpyridinium iodide and stirring was continued for 30 minutes at room temperature. Then was added a solution of 3.68 parts of 4-[[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-ylamino]-1-piperidine]ethanol in 130 parts of dichloromethane and the whole was stirred for 1 hour at room temperature. The reaction mixture was washed with water. The organic phase was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (96:4 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (E)-2-butenedioate salt in methanol. The salt was filtered off and dried, yielding 0.3 parts (4%) of [2-4-[[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-2-oxo-2*H*-benzopyran-3-carboxylate (E)-2-butanedioate (1:2); mp. 205.0°C (compound 42).

25 Following the same procedure and using equivalent amounts of the appropriate starting materials, there was also prepared:

N-[2-4-[[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-4-oxo-4*H*-benzopyran-2-carboxamide (E)-2-butenedioate (1:2); mp. 248.7°C (compound 43).

Example 33

30 To a stirred and cooled (below 10°C) mixture of 3.8 parts of 2-oxo-2*H*-1-benzopyran-3-carboxylic acid, 2.2 parts of *N,N*-diethylethanamine and 225 parts of trichloromethane was added dropwise a solution of 1.9 parts of methyl carbonochloride in 75 parts of trichloromethane. Upon completion, stirring was continued for 30 minutes at room temperature. This solution was added dropwise to a stirred and cooled solution of 6.6 parts of *N*-[1-(2-aminoethyl)-4-piperidinyl]-1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-amine in 75 parts of trichloromethane at 5°C. The whole was stirred for 1 hour while the mixture was allowed to reach room temperature. The reaction mixture was washed successively with water, a sodium hydroxide solution in 10% and again with water. The organic phase was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (E)-2-butenedioate salt in methanol. The salt was filtered off and dried, yielding 6.8 parts (47.5%) of *N*-[2-4-[[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-2-oxo-2*H*-1-benzopyran-3-carboxamide (E)-2-butenedioate (1:2); mp. 216.8°C (compound 44).

Example 34

45 A mixture of 4.4 parts of *N*-(5-bromo-1,3,4-thiadiazol-2-yl)-acetamide, 7.3 parts of *N*-[1-(2-aminoethyl)-4-piperidinyl]-1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-amine, 3.18 parts of sodium carbonate and 135 parts of *N,N*-dimethylformamide was stirred overnight at 80–90°C. The reaction mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from a mixture of acetonitrile and 2,2'-oxybispropane, yielding 1.7 parts of *N*-[2-4-[[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-formamide; mp. 153.2°C (compound 45).

Example 35

50 A mixture of 5.09 parts of *N*-[1-(2-aminoethyl)-4-piperidinyl]-1-(2-furanyl)methyl]-1*H*-benzimidazol-2-amine and 54 parts of *N,N*-dimethylformamide was stirred and heated at 50°C and there was added dropwise a solution of 2.8 parts of dihydro-3-phenyl-2*H*-pyran-2,6(3*H*)dione in 18 parts of *N,N*-dimethylformamide. Upon completion, stirring was continued overnight at 50°C. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (95:5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was from a mixture of acetonitrile and 2,2'-oxybispropane, yielding 1.8 parts of *N*-[2-4-[[1-[(2-furanyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]-formamide hemihydrate; mp. 125.2°C (compound 46).

Example 36

A mixture of 30 parts of 2-furanmethanol, 300 parts of a formaldehyde solution 4% in water and 145 parts of 1-[(4-fluorophenyl)methyl]-*N*-(4-piperidinyl)-1*H*-benzimidazol-2-amine dihydrobromide was stirred at 3°C. The mixture was allowed to reach slowly room temperature and stirring was continued for 3 days at room temperature. The reaction mixture was alkalinized and extracted with dichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using first trichloromethane and then a mixture of trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (E)-2-butenedioate salt and the free base was liberated again in the conventional manner. It was crystallized from a mixture of 2-propanone and 2,2'-oxybispropane, yielding 57 parts (44%) of 5-[4-[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]methyl]-2-furanmethanol; mp. 148.8°C (compound 47).

Example 37

To a stirred solution of 6.5 parts of 5-[4-[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]methyl]-2-furanmethanol in 180 parts of *N,N*-dimethylformamide was added portionwise 1 part of a sodium hydride dispersion 50% at room temperature. After stirring for 1 hour, a solution of 1.6 parts of 2-chloropyrimidine in *N,N*-dimethylformamide was added dropwise. Upon completion, stirring was continued overnight at room temperature. The reaction mixture was poured into water and the product was extracted with dichloromethane. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using first trichloromethane and then a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The first fraction was collected and the eluent was evaporated. The residue was crystallized from a mixture of 4-methyl-2-pentanone and 2,2'-oxybispropane, yielding 2.1 parts of 1-[(4-fluorophenyl)methyl]-*N*-[1-[5-[2-(pyrimidin-2-yl)oxy]methyl]-2-furanyl]methyl]-4-piperidinyl]-1*H*-benzimidazol-2-amine; mp. 167.8°C (compound 48).

Example 38

To a stirred solution (0°C) of 11.4 parts of 2-aminoethanethiol hydrochloride in 48 parts of concentrate hydrochloric acid were added portionwise 25 parts of 5-[4-[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]methyl]-2-furanmethanol. Upon completion, stirring was continued first overnight at 0°C and then for 4 days at room temperature. The reaction mixture was alkalinized with a dilute potassium hydroxide solution and the product was extracted with dichloromethane. The extract was dried, filtered and evaporated. The residue was purified by filtration over silica gel using a mixture of trichloromethane and methanol, saturated with ammonia, (90:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated, yielding 24 parts (88.5%) of *N*-[1-[5-[2-(2-aminoethylthiomethyl)-2-furanyl]methyl]-4-piperidinyl]-1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-amine as an oily residue (49).

A mixture of 1.14 parts of 2-chloropyrimidine, 5 parts of *N*-[1-[5-[2-(2-aminoethylthiomethyl)-2-furanyl]methyl]-4-piperidinyl]-1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-amine, 8 parts of sodium hydrogen carbonate and 80 parts of ethanol was stirred and refluxed overnight. The reaction mixture was filtered and the filtrate was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (97:3 by volume) as eluent. The main fraction was collected and the eluent was evaporated. The residue was crystallized from 1,1'-oxybisethane, yielding 1.2 parts (21%) of 1-[(4-fluorophenyl)methyl]-*N*-[1-[5-[2-(2-pyrimidinylamino)ethyl]thiomethyl]-2-furanyl]methyl]-4-piperidinyl]-1*H*-benzimidazol-2-amine; mp. 128.9°C (compound 50).

Example 39

A mixture of 7.7 parts of 2-amino-*N*-[2-[4-[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]benzamide, 20 parts of acetic anhydride and 80 parts of water was stirred for 4 hours at 100°C. Water was added and the whole was alkalinized with ammonium hydroxide. The product was extracted with trichloromethane. The extract was dried, filtered and evaporated. The residue was crystallized from 4-methyl-2-pentanone. The product was filtered off and recrystallized from acetonitrile, yielding 7.7 parts of 2-(acetylaminio)-*N*-[2-[4-[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]benzamide; mp. 170.9°C (compound 51).

In a similar manner there was also prepared:

2-(acetylaminio)-*N*-[2-[4-[1-[(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]benzamide (E)-2-butenedioate (1:2); mp. 175.2°C (compound 52).

Example 40

To a stirred mixture of 4.4 parts of *N*-[1-[2-(4-aminophenyl)ethyl]-4-piperidinyl]-1-(4-fluorophenyl)methyl]-1*H*-benzimidazol-2-amine, 1.05 parts of *N,N*-diethylethanamine and 195 parts of dichloromethane were added dropwise 1.14 parts of methanesulfonyl chloride. Upon completion, stirring was continued for 3 hours at room temperature. Water was added and the whole was alkalinized with a sodium hydroxide solution. The organic phase was separated, dried, filtered and evaporated. The residue

was separated by HPLC over silica gel using a mixture of trichloromethane, hexane and methanol (45:45:10 by volume) as eluent. The first fraction was collected and the eluent was evaporated. The residue was crystallized from acetonitrile, yielding 1.8 parts of *N*-[4-[2-[4-[[1-[4-fluorophenyl]methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]methanesulfonamide monohydrate; mp. 191.0°C (compound 53).

In a similar manner there were also prepared:

N-[4-[2-[4-[[1-[4-fluorophenyl]methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]phenyl]benzamide monohydrochloride; mp. 217.3°C (compound 54).

N-[4-[2-[4-[[1-[4-fluorophenyl]methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]phenyl]acetamide; mp. 227.2°C (compound 55).

Example 41

To a stirred mixture of 4.4 parts of *N*-[1-[2-(4-aminophenyl)ethyl]-4-piperidinyl]-1-[4-fluorophenyl]methyl]-1*H*-benzimidazol-2-amine, 16 parts of acetic acid and 32 parts of water was added dropwise a solution of 1.2 parts of potassium isocyanate in 33 parts of water. Upon completion, stirring was continued overnight at room temperature. The reaction mixture was evaporated and the residue was taken up in water and dichloromethane. The whole was alkalinized with sodium hydroxide. The precipitated product was filtered off and purified by HPLC over silica gel using a mixture of trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from acetonitrile, yielding 1.3 parts of *N*-[4-[2-[4-[[1-[4-fluorophenyl]methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]phenyl]urea; mp. 186.9°C (compound 56).

Example 42

A mixture of 0.6 parts of isocyanatomethane, 4.43 parts of *N*-[1-[2-(4-aminophenyl)ethyl]-4-piperidinyl]-1-[4-fluorophenyl]methyl]-1*H*-benzimidazol-2-amine and 135 parts of tetrahydrofuran was stirred for 3 hours at room temperature. The precipitated product was filtered off and crystallized from acetonitrile, yielding 2 parts (39.9%) of *N*-[4-[2-[4-[[1-[4-fluorophenyl]methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]phenyl]-*N'*methylurea; mp. +300°C (compound 57).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there was also prepared:

N-[4-[2-[4-[[1-[4-fluorophenyl]methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]phenyl]-*N'*-methylthiourea monohydrate; mp. 120.2°C (compound 58).

Example 43

A mixture of 5 parts of *N*-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]-1-[4-thiazolyl]methyl]-1*H*-benzimidazol-2-amine and 150 parts of a hydrobromic acid solution 48% in water was stirred and refluxed overnight. The reaction mixture was evaporated and the solid residue was crystallized from ethanol 80%, yielding 4 parts of 4-[2-[4-[[1-[4-thiazolyl]methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]phenol dihydrobromide; mp. 291.0°C (compound 59).

Example 44

A mixture of 5 parts of *N'*-cyano-*N*-(2,2-dimethoxyethyl)-*N'*-[2-[4-[[1-[4-fluorophenyl]methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]guanidine and 60 parts of concentrated hydrochloric acid was stirred and refluxed for one hour. Water was added and the whole was alkalinized with ammonia. The product was extracted with 4-methyl-2-pentanone. The extract was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) saturated with ammonia, as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from a mixture of acetonitrile and 2,2'-oxybispropane, yielding 1 part of *N*-[1-[2-(2-amino-1*H*-imidazol-1-yl)ethyl]-4-piperidinyl]-1-[4-fluorophenyl]methyl]-1*H*-benzimidazol-2-amine monohydrate; mp. 171.4°C (compound 60).

C. Formulations

The following formulations exemplify typical pharmaceutical compositions in dosage unit form suitable for systemic administration to animal and human subjects in accordance with the present invention. These examples are given to illustrate the scope of the present invention.

"Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I), a possible stereochemically isomeric form or pharmaceutically acceptable acid addition salt thereof.

Example 45

Oral drops

500 Grams of the A.I. was dissolved in 0.5 liters of 2-hydroxy-propanoic acid and 1.5 liters of the polyethylene glycol at 60–80°C. After cooling to 30–40°C there were added 35 liters of polyethylene glycol and the mixture was stirred well. Then were added a solution of 1750 grams of sodium saccharin in 2.5 liters of purified water and while stirring there were added 2.5 liters of cocoa flavor and polyethylene glycol q.s. to a volume of 50 liters, providing an oral drop solution comprising 10 milligrams of the A.I. per

EP 0 151 824 B1

milliliter. The resulting solution was filled into suitable containers.

Example 46

Oral solution

9 Grams of methyl 4-hydroxybenzoate and 1 gram of propyl 4-hydroxybenzoate were dissolved in 4 liters of boiling purified water. In 3 liters of this solution were dissolved first 10 grams of 2,3-dihydroxybutanedioic acid and thereafter 20 grams of the A.I. The latter solution was combined with the remaining part of the former solution and 12 liters 1,2,3-propanetriol and 3 liters of sorbitol 70% solution were added thereto. 40 Grams of sodium saccharin were dissolved in 0.5 liters of water and 2 milliliters of raspberry and 2 milliliters of gooseberry essence were added. The latter solution was combined with the former, water was added q.s. to a volume of 20 liters providing an oral solution comprising 20 milligrams of the active ingredient per teaspoonful (5 milliliters). The resulting solution was filled in suitable containers.

Example 47

Capsules

20 Grams of A.I., 6 grams sodium lauryl sulfate, 56 grams starch, 56 grams lactose, 0.8 grams colloidal silicon dioxide, and 1.2 grams magnesium stearate were vigorously stirred together. The resulting mixture was subsequently filled into 1000 suitable hardened gelating capsules, comprising each 20 milligrams of the active ingredient.

Film-coated tablets

Preparation of tablet core

A mixture of 100 grams of the A.I., 570 grams lactose and 200 grams starch was mixed well and thereafter humidified with a solution of 5 grams sodium dodecyl sulfate and 10 grams polyvinylpyrrolidone in about 200 milliliters of water. The wet powder mixture was sieved, dried and sieved again. Then there was added 100 grams microcrystalline cellulose and 15 grams hydrogenated vegetable oil. The whole was mixed well and compressed into tablets, giving 10,000 tablets, each containing 10 milligrams of the active ingredient.

Coating

To a solution of 10 grams methyl cellulose in 75 milliliters of denaturated ethanol there was added a solution of 5 grams of ethyl cellulose in 150 milliliters of dichloromethane. Then there were added 75 milliliters of dichloromethane and 2.5 milliliters 1,2,3-propanetriol. 10 Grams of polyethylene glycol was molten and dissolved in 75 milliliters of dichloromethane. The latter solution was added to the former and then there were added 2.5 grams of magnesium octadecanoate, 5 grams of polyvinylpyrrolidone and 30 milliliters of concentrated colour suspension (Opaspray K—1—2109) and the whole was homogenated.

The tablet cores were coated with the thus obtained mixture in a coating apparatus.

Example 49

Injectable solution

1.8 Grams methyl 4-hydroxybenzoate and 0.2 grams propyl 4-hydroxybenzoate were dissolved in about 0.5 liters of boiling water for injection. After cooling to about 50°C there were added while stirring 4 grams lactic acid, 0.05 propylene glycol and 4 grams of the A.I. The solution was cooled to room temperature and supplemented with water for injection q.s. ad 1 liter volume, giving a solution of 4 milligrams A.I. per milliliters. The solution was sterilized by filtration (U.S.P. XVII p. 811) and filled in sterile containers.


Example 50

Suppositories

3 Grams A.I. was dissolved in a solution of 3 grams 2,3-dihydroxybutanedioic acid in 25 milliliters polyethylene glycol 400. 12 Grams surfactant and triglyceride q.s. ad 300 grams were molten together. The latter mixture was mixed well with the former solution. The thus obtained mixture was poured onto moulds at a temperature of 37—38°C to form 100 suppositories each containing 30 milligrams of the active ingredient.

Claims

1. A chemical compound having the formula



(I),

a pharmaceutically acceptable acid addition salt or a possible stereochemically isomeric form thereof, wherein:

$A^1=A^2-A^3=A^4$ is a bivalent radical having the formula




wherein one or two hydrogen atoms in said radicals (a-1) – (a-5) may, each independently from each other, be replaced by halo, lower alkyl, lower alkyloxy, trifluoromethyl or hydroxy;

R is a member selected from the group consisting of hydrogen and lower alkyl;

R¹ is a member selected from the group consisting of hydrogen, alkyl, cycloalkyl, Ar¹ and lower alkyl substituted with one or two Ar¹ radicals.

R^2 is a member selected from the group consisting of hydrogen, lower alkyl, cycloalkyl, (lower alkyl)-CO-, (lower alkyloxy)-CO- and Ar^2 -lower alkyl; and

L is a radical of formula



i) where $A^1=A^2-A^3=A^4$ is a radical of formula (a-3), (a-4) or (a-5), or

ii) where $A^1=A^2-A^3=A^4$ is a radical of formula (a-1) or (a-2), and R^1 is Ar^3 or lower alkyl substituted with one or two Ar^3 radicals, said Ar^3 being pyrazinyl, thiazolyl or imidazolyl, optionally substituted with lower alkyl;


L may also be a radical of formula:



said W being a member selected from the group consisting of hydrogen, lower alkyl, Ar^1 , Ar^1 -lower alkyl, 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl, a radical of formula



a radical of formula



(c-1-b), or

a radical of formula



(c-1-c)

wherein R^3 and R^4 each independently hydrogen or lower alkyl; and W^1 is cycloalkyl or lower alkyl, optionally substituted with up to two substituents selected from the group consisting of hydroxy, lower alkyloxy, 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl and Ar^1 ; and where Z^1 is NR^8 , W^1 may also be hydrogen, amino, lower alkylamino, Ar^1 -amino or nitro;


said W^2 being a member selected from the group consisting of hydrogen, lower alkyl, Ar^1 and a radical of formula:



(c-2-a)

wherein R^8 is hydrogen, lower alkyl or Ar^1 ;

said T being a radical of formula:



(c-3-a),


(c-3-b);

R^6 being hydrogen, lower alkyl or Ar^1 ;

R^7 being lower alkyl or Ar^1 ; and


R^8 being hydrogen or lower alkyl;

said Het being a radical of formula (c-1-a), (c-1-b), or a radical of formula




(c-4-a)

or of formula



(c-4-b),


wherein R^9 , R^{10} , R^{11} and R^{12} are each independently hydrogen or lower alkyl; or a radical of formula



(c-4-c),

wherein R^{13} is hydrogen, lower alkyl or amino, or

said Het being furan substituted with lower alkyl, said lower alkyl being optionally substituted with hydroxy, mercapto, lower alkyloxy, lower alkylthio, (aminolower alkyl)thio, Ar^1-O- or with a radical of formula



(c-4-d-1),

s being an integer of from 1 to 6 inclusive; or where Z or Y is a direct bond, s may also be 0; and R^{14} being hydrogen or lower alkyl; wherein

n is 0 or the integer of 1 or 2;

X is O, S, NR^{15} or $CHNO_2$;

Y is O, S, NR^{16} or a direct bond;

Y^1 is O, S or NR^{16} ;

Y^2 is S or NR^{16} ;

Z is O, S, NR^8 or a direct bond;

Z¹ is O, S or NR⁸;

X^a and Y^a independently having the same meaning of X respectively Y;

said R^{1b} being hydrogen, lower alkyl, cyano, nitro, Ar²-sulfonyl, lower alkylsulfonyl, lower alkylcarbonyl or Ar²-carbonyl;

said R^{1b} being hydrogen, lower alkyl, (Ar²)lower alkyl, 2-lower alkyloxy-1,2-dioxoethyl; or a radical of formula —C(=X)—R^{17} ; R¹⁷ being hydrogen, lower alkyl, Ar², Ar²-lower alkyl, lower alkyloxy, Ar²-lower alkyloxy, mono- or di(lower alkyl)amino, Ar²-lower alkylamino or Ar²-lower alkyl(lower alkyl)amino;

wherein Ar¹ is a member selected from the group consisting of phenyl, being optionally substituted with up to three substituents each independently selected from the group consisting of halo, hydroxy, nitro, cyano, trifluoromethyl, lower alkyl, lower alkyloxy, mercapto, amino, mono- and di(lower alkyl)amino, carboxyl, lower alkyloxy carbonyl and lower alkyl-CO—; thienyl; halothienyl; furanyl; lower alkyl substituted furanyl; pyridinyl; pyrazinyl; thiazolyl and imidazolyl optionally substituted with lower alkyl; and wherein Ar² is a member selected from the group consisting of phenyl being optionally substituted with up to three substituents each independently selected from the group consisting of halo, hydroxy, nitro, cyano, trifluoromethyl, lower alkyl, lower alkyloxy, lower alkylthio, mercapto, amino, mono- and di(lower alkyl)amino, carboxyl, lower alkyloxy carbonyl and lower alkyl-CO—; wherein lower alkyl represents straight and branched-chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms; alkyl includes lower alkyl and the higher homologs thereof having from 7 to 10 carbon atoms; cycloalkyl is cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; lower alkenyl represents a straight or branched-chain hydrocarbon radical containing one double bond and having 2 to 6 carbon atoms; lower alkanediy l represents bivalent, straight or branched-chain alkanediy l radicals having from 1 to 6 carbon atoms;

provided that:

i) When A¹=A²—A³=A⁴ is a radical of formula (a-1) or (a-2), and L is a radical of formula (b-1), wherein W is other than hydrogen or other than a radical of formula (c-1-a) or (c-1-b), then X is other than O;

ii) when L is a radical of formula (b-1), wherein W is a radical of formula (c-1-c), wherein Z¹ is NH then W¹ is other than hydrogen or lower alkyl;

iii) when A¹=A²—A³=A⁴ is a radical of formula (a-1) or (a-2), and L is a radical of formula (b-3), wherein X is O, Y is NR^{1b}, O or a direct bond, and X^a is O,

a) then Y^a is not O;

b) and W² being lower alkyl then Y^a is not a direct bond;

2. A chemical compound according to claim 1, wherein L is a radical of formula (b-1), wherein Y is NH, X is O and W is hydrogen; or L is a radical of formula (b-1) wherein X is S, NH or NCN, Y is NH and W is 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl, or a radical of formula (c-1-c), wherein Z¹ is NR⁸ and W¹ is amino, nitro or lower alkyl, optionally substituted with one hydroxy, lower alkyloxy, 1-piperidinyl, 1-pyrrolidinyl, 4-morpholinyl or phenyl radical, or with two lower alkyloxy radicals; or L is a radical of formula (b-1), wherein X is S, NH or NCN, Y is NH and W is lower alkyloxy or lower alkylthio; or wherein L is a radical of formula (b-1) wherein W is a radical of formula (c-1-a) or (c-1-b); or

L is a radical of formula (b-2) wherein n is 1, X is O or S and W is a radical of formula (c-1-c), wherein Z¹ is NR⁸ and W¹ is lower alkyl; or

L is a radical of formula (b-3), wherein X is O, Y is NH, X^a is O, Y^a is NR^{1b} and W² is lower alkyl; or

L is a radical of formula (b-4), wherein T is a radical of formula (c-3-a), wherein X is O or S, Z is NR⁸ and R⁸ is hydrogen or lower alkyl; or wherein T is a radical of formula (c-3-b), wherein R⁸ is hydrogen and R⁷ is lower alkyl; or

L is a radical of formula (b-5) wherein Het is a radical of formula (c-4-a), wherein R⁸, R¹¹ and R¹² are hydrogen; or wherein Het is a radical of formula (c-4-c); or wherein Het is furan substituted with lower alkyl being substituted with hydroxy or with a radical of formula (c-4-d-1), wherein Y is O or S, Z is NH or a direct bond and R¹⁴ is hydrogen; or

L is a radical of formula (b-6) wherein Y¹ is O; or

L is a radical of formula (b-7) wherein Ar¹ is phenyl substituted with hydroxy or lower alkyloxy.

3. A chemical compound according to any of claims 1 and 2 for use as a medicine.

4. A chemical compound according to any of claims 1 and 2 for use as an anti-allergic medicine.

5. A pharmaceutical composition comprising an inert carrier and a pharmaceutically acceptable amount of a compound according to any of claims 1 and 2.

6. A pharmaceutical composition according to claim 5 for use as an anti-allergic medicine.

7. A process for preparing a pharmaceutical composition, characterized in that a therapeutically effective amount of a compound as claimed in any of claims 1 and 2 is intimately mixed with a suitable pharmaceutical carrier.

8. A process for preparing a compound according to any of claims 1 and 2 characterized by

a) alkylating a piperidine of formula Q²—D (III) with an intermediate of formula Q¹ (II) in a reaction-inert solvent wherein


1) Q² is hydrogen and Q¹ is an intermediate of formula L—G (II-a), said G representing halo or sulfonyloxy; or

2) Q¹ is an intermediate of formula W—C(X)=G¹, (II-b-1), said G¹ representing halo, sulfonyloxy, lower alkyloxy, lower alkylthio, Ar²-oxy, Ar²-thio, lower alkylcarbonyloxy or lower alkyloxy carbonyloxy, or


where G¹ is connected to




it may also be —N(loweralkyl)NO; and Q² is a radical of formula HY¹—Alk—, thus preparing a compound of formula



3) Q¹ is an intermediate of formula




and Q² is a radical of formula HY¹—Alk—, thus preparing a compound of formula



4) Q¹ is an intermediate of formula (lower alkenyl)—G, (II-b-3), and Q² is a radical HY¹—Alk—, thus preparing a compound of formula (lower alkenyl)—Y¹—Alk—D (I-a-3); or

5) Q¹ is an intermediate of formula



and Q² is a radical of formula



thus preparing a compound of formula



6) Q¹ is an intermediate of formula W—C(=X)—Y¹H (II-c-1) and Q² is a radical of formula G—Alk, thus preparing a compound of formula (I-a-1);

7) Q¹ is an intermediate of formula



and Q² is a radical of formula G—Alk, thus preparing a compound of formula (I-a-2);

8) Q¹ is an intermediate of formula (lower alkenyl)—Y¹—H, (II-c-3) and Q² is a radical of formula G—Alk—, thus preparing a compound of formula (I-a-3);

9) Q¹ is an intermediate of formula Het¹—H, (II-c-4), wherein said Het¹ is a radical of formula (c-4-a), (c-4-b) or (c-4-c), and Q² is a radical of formula G—Alk, thus preparing a compound of formula Het¹—Alk—D (I-a-5); or

b) reacting a reagent of formula W¹—Z¹H (II-d) with an intermediate of formula HY¹—Alk—D (III-b-1-) in

EP 0 151 824 B1

the presence of an appropriate $C=X$ generating agent, in a reaction-inert solvent, thus preparing a compound of formula $W^1Z^1-C(=X)-Y^1-Alk-D$, (I-a-1-a); or
c) reacting a reagent of formula (II-d) with an intermediate of formula




in the presence of an appropriate



generating agent, in a reaction-inert solvent, thus preparing a compound of formula




d) cyclodesulfurizing an intermediate of formula



with an appropriate alkyl halide, metal oxide or metal salt in a reaction-inert solvent; or

e) reacting an intermediate of formula W^1-Z^1-H (V) with a piperidine of formula $X^1=C=N-Alk-D$ (III-b-1), wherein X^1 is O or S, in a reaction-inert solvent, thus preparing a compound of formula



f) reacting an intermediate of formula $W^1-N=C=X^1$ (VII), with a piperidine of formula $HY^1-Alk-D$ (III-b-1) in a reaction-inert solvent, thus preparing a compound of formula



g) reacting an intermediate of formula (VII) with a piperidine of formula




in a reaction-inert solvent, thus preparing a compound of formula



h) reacting an intermediate of formula $W^2-C(=X^1)-OH$ (VIII), said W^2 having the previously described meaning of W , provided that W^2 is other than a radical of formula (c-1-c), with a piperidine of formula $HY^1-Alk-D$ (III-b-1) in a reaction-inert solvent, if desired, after converting the OH-function in (VIII) in a suitable leaving group, or, if desired, by reacting (III-b-1) with (VIII) together with an appropriate reagent capable of forming amides or esters, thus preparing a compound of formula

EP 0 151 824 B1



i) reacting an intermediate of formula $W^3-C(=X^1)-OH$ (VIII), said W^3 having the previously defined meaning, with a piperidine of formula



in a reaction-inert solvent, if desired, after converting the OH-function in (VIII) in a suitable leaving group, or, if desired, by reacting (III-2-b) with (VIII) together with an appropriate reagent capable of forming amides or esters, thus preparing a compound of formula



j) reacting a piperidine of formula HD (III-a) with a reagent of formula L^1 -lower alkenediyl-H (IX), wherein L^1 is selected so that it forms, combined with -Alk-, a radical of formula (b-1), (b-3), (b-4), (b-5), (b-6) or (b-7), in a suitable reaction-inert solvent, thus preparing a compound of formula $L^1-Alk-D$ (I-d); or

k) reacting a furan of formula




with a piperidine of formula (III-a) in the presence of formaldehyde or a polymeric form thereof, in a reaction-inert solvent, thus preparing a compound of



wherein R^{18} is an in claim 1 defined substituent of said furan ring; or


l) alkylating a furan derivative of formula $H-Y^1-D^1$, (X-a), wherein D^1 represents a radical of formula



with an intermediate of formula $E-Z-C_6H_{28}-G$, (XI-a), wherein E represents a radical of formula



in reaction-inert solvent, thus preparing a compound of formula




m) alkylating an intermediate of formula $E-Z-C_6H_{28}-Y^1H$, (XI-b), with a furan derivative of formula $G-D^1$, (X-b), in a reaction-inert solvent, thus preparing a compound of formula (I-e-1);

n) alkylating a furan derivative of formula $H-Z^1-C_6H_{28}-Y-D^1$ (X-c), with an intermediate of formula $E-G$, (XI-c), in a reaction-inert solvent, thus preparing a compound of formula




c) alkylating an intermediate of formula $E-Z^1H$, (XI-d), with a furan derivative of formula $G-C_6H_{28}-Y-D^1$, (X-d), in a reaction-inert solvent, thus preparing a compound of formula (I-e-2);


p) reacting an intermediate of formula




wherein $\text{Y}^{\text{a-1}}$ has the previously described meaning of Y^{a} , provided that it is not a direct bond, with a reagent of formula $\text{W}^{\text{a}}-\text{C}(\text{X}^{\text{a}})=\text{G}^1$, (XIII-a), in a reaction-inert solvent, thus preparing a compound of formula




q) reacting an intermediate of formula




with a reagent of formula $\text{R}^{\text{e}}-\text{Z}-\text{C}(\text{X})=\text{G}^1$, in a reaction-inert solvent, thus preparing a compound of formula




r) reacting an intermediate of formula




with a reagent of formula $\text{R}^7-\text{SO}_2-\text{G}^1$, in a reaction-inert solvent, thus preparing a compound of formula




s) reacting a reagent of formula $\text{R}^{\text{e}}-\text{N}=\text{C}=\text{X}^{\text{a-1}}$ (XIV-a), wherein $\text{X}^{\text{a-1}}$ is O or S, with an intermediate of formula (XII-a) in a reaction-inert solvent, thus preparing a compound of formula



t) reacting an intermediate of formula (XII-b) with a reagent of formula $\text{R}^{\text{e}}-\text{N}=\text{C}=\text{X}^1$ (XIV-b), thus preparing a compound of formula




45 wherein D represents a radical of formula



10 optionally converting the compounds of formula (I) into each other following art-known functional group transformation procedures; and, if desired, converting the compounds of formula (I) into a therapeutically active non-toxic acid-addition salt form by treatment with an appropriate acid or, conversely, converting the acid-addition salt into the free base form with alkali; and/or preparing stereochemically isomeric forms thereof.

Patentansprüche

1. Chemische Verbindung der Formel:



ein pharmazeutisch annehmbares Säureadditionssalz oder eine mögliche stereochemisch isomere Form hiervon, worin:

$$A^1 = A^2 - A^3 = A^4 \text{ ein zweiwertiger Rest mit der Formel}$$




40 ist, worin ein oder zwei Wassersatoffatome in den genannten Resten (a-1) bis (a-5) jeweils unabhängig voneinander durch Halogen, Niederalkyl, Niederalkyloxy, Trifluormethyl oder Hydroxy ersetzt sein können:

R ein Vertreter, ausgewählt aus der aus Wasserstoff und Niederalkyl bestehenden Gruppe, ist;

R¹ ein Vertreter, ausgewählt aus der aus Wasserstoff, Alkyl, Cycloalkyl, Ar¹ und Niederalkyl, substituiert mit einem oder zwei Ar¹-Resten, bestehenden Gruppe, ist;

45 R^2 ein Vertreter, ausgewählt aus der aus Wasserstoff, Niederkalkyl, Cycloalkyl, (Niederkalkyl)-CO-, (Niederkalkyloxy)-CO- und Ar^2 -Niederkalkyl bestehenden Gruppe, ist; und

L ein Rest der Formel



Het-Alk-

(b-5);

(Niederalkenyl)-Y¹-Alk-

(b-6)

ist, oder


i) wenn A¹=A²-A³=A⁴ ein Rest der Formel (a-3), (a-4) oder (a-5) ist, oderii) wenn A¹=A²-A³=A⁴ ein Rest der Formel (a-1) oder (a-2) ist, und R¹ Ar¹ oder Niederalkyl, substituiert mit einem oder zwei Ar²-Resten, bedeutet, welches Ar³ Pyrazinyl, Thiazolyl oder Imidazolyl, wahlweise substituiert mit Niederalkyl, bedeutet,

L auch ein Rest der Formel

Ar¹-Alk-


(b-7)

sein kann; worin

W ein Vertreter, ausgewählt aus der aus Wasserstoff, Niederalkyl, Ar¹, Ar¹-Niederalkyl, 1-Piperidiny, 1-Pyrrolidiny, 4-Morpholiny, einem Rest der Formel

(c-1-a),

einem Rest der Formel



(c-1-b), oder

einem Rest der Formel

W¹-Z¹-

(c-1-c)

bestehenden Gruppe ist, worin

R³ und R⁴ voneinander unabhängig Wasserstoff oder Niederalkyl sind; undW¹ für Cycloalkyl oder Niederalkyl, wahlweise substituiert mit bis zu 2 Substituenten, ausgewählt aus der aus Hydroxy, Niederalkyloxy, 1-Piperidiny, 1-Pyrrolidiny, 4-Morpholiny und Ar¹ bestehenden Gruppe, ist; und wennZ¹ für NR⁸ steht, W¹ auch Wasserstoff, Amino, Niederalkylamino, Ar¹-Amino oder Nitro sein kann;

worin


W² ein Vertreter aus der aus Wasserstoff, Niederalkyl, Ar¹ und einem Rest der Formel:R⁵-Z¹-

(c-2-a)

bestehenden Gruppe ist, worin

R⁵ für Wasserstoff, Niederalkyl oder Ar¹ steht; worin

T ein Rest der Formel:




(c-3-a),

(c-3-b)

ist; wobei


R⁶ Wasserstoff, Niederalkyl oder Ar¹ ist; wobeiR⁷ Niederalkyl oder Ar¹ ist; und wobeiR⁸ Wasserstoff oder Niederalkyl ist; worin

Het ein Rest der Formel (c-1-a), (c-1-b) oder ein Rest der Formel




(c-4-a)

oder der Formel



(c-4-b),

ist, worin R⁹, R¹⁰, R¹¹ und R¹² unabhängig voneinander Wasserstoff oder Niederalkyl sind; oder ein Rest der Formel




(c-4-c),

ist, worin

R¹³ für Wasserstoff, Niederalkyl oder Amino steht, oder worin

Het Furan, substituiert mit Niederalkyl, ist, worin Niederalkyl gegebenenfalls mit Hydroxy, Mercapto, Niederalkylthio, Niederalkylthio, (Aminoniederalkyl)thio, Ar'-O- oder mit einem Rest der Formel



(c-4-d-1),

substituiert ist, worin

s eine ganze Zahl von 1 bis einschließlich 6 ist; oder wenn Z oder Y eine direkte Bindung darstellen, s auch 0 sein kann; und R¹⁴ Wasserstoff oder Niederalkyl ist; worin

n den Wert 0 besitzt oder die ganze Zahl 1 oder 2 bedeutet;

X für O, S, NR¹⁶ oder CHNO₂ steht;

Y für O, S, NR¹⁶ oder eine direkte Bindung steht;

Y¹ für O, S oder NR¹⁶ steht;

Y² S oder NR¹⁶ ist;

Z für O, S, NR¹⁶ oder eine direkte Bindung steht;

Z¹ für O, S oder NR¹⁶ steht;

X^a und Y^a unabhängig voneinander die gleiche Bedeutung von X bzw. Y besitzen; welches

R¹⁶ Wasserstoff, Niederalkyl, Cyano, Nitro, Ar²-Sulfonyl, Niederalkylsulfonyl, Niederalkylcarbonyl oder

Ar²-Carbonyl ist; welches

R¹⁶ Wasserstoff, Niederalkyl, (Ar²)Niederalkyl, 2-Niederalkyl-1,2-dioxoethyl oder ein Rest der

Formel —C(=X)—R¹⁷ ist; worin R¹⁷ Wasserstoff, Niederalkyl, Ar², Ar²-Niederalkyl, Niederalkylthio, Ar²-

Niederalkylthio, Mono- oder Di(niederalkyl)amino, Ar²-Niederalkylamino oder Ar²-Niederalkyl(niederal-

kyl)amino ist; worin

Ar² ein Vertreter, ausgewählt aus der aus Phenyl, das gegebenenfalls mit bis zu 3 Substituenten

substituiert ist, welche voneinander unabhängig aus der aus Halogen, Hydroxy, Nitro, Cyano, Trifluormethyl, Niederalkyl, Niederalkylthio, Niederalkylthio, Mercapto, Amino, Mono- und

Di(niederalkyl)amino, Carboxyl, Niederalkylcarbonyl und Niederalkyl—CO— bestehenden Gruppe

ausgewählt sind; Thienyl; Halogenthienyl; Furanyl; Niederalkyl-substituiertes Furanyl; Pyridinyl; Pyrazinyl; Thiazolyl und Imidazolyl, das gegebenenfalls mit Niederalkyl substituiert ist, bestehenden

Gruppe ist; und worin

Ar² ein Vertreter ausgewählt aus der Gruppe bestehend aus Phenyl, das gegebenenfalls mit bis zu 3

Substituenten substituiert ist, welche voneinander unabhängig aus der aus Halogen, Hydroxy, Nitro, Cyano, Trifluormethyl, Niederalkyl, Niederalkylthio, Niederalkylthio, Mercapto, Amino, Mono- und

Di(niederalkyl)amino, Carboxyl, Niederalkylcarbonyl und (Niederalkyl)-CO bestehenden Gruppe

ausgewählt sind, ist; worin

Niederalkyl lineare und verzweigte gesättigte Kohlenwasserstoffreste mit 1 bis 6 Kohlenstoffatomen

bedeutet; Alkyl, Niederalkyl und die höheren Homologen hiervon mit 7 bis 10 Kohlenstoffatomen umfaßt;

Cycloalkyl für Cyclopropyl, Cyclobutyl, Cyclopentyl und Cyclohexyl steht; Niederalkyl einen linearen

oder verzweigten Kohlenwasserstoffrest mit einer Doppelbindung und 2 bis 6 Kohlenstoffatomen

darstellt; Niederalcandyl zweiwertige lineare oder verzweigte Alkandylreste mit 1 bis 6

Kohlenstoffatomen bedeutet; mit der Maßgabe daß:

i) wenn A¹=A²=A³=A⁴ ein Rest der Formel (a-1) oder (a-2) ist und L einen Rest der Formel (b-1)

bedeutet, worin W eine andere Bedeutung als Wasserstoff oder eine andere Bedeutung als die eines Restes

der Formel (c-1-a) oder (c-1-b) besitzt, X dann eine andere Bedeutung als O besitzt;

ii) wenn L ein Rest der Formel (b-1) ist, worin W ein Rest der Formel (c-1-c) ist, worin Z¹ für NH steht, W¹

dann eine andere Bedeutung als Wasserstoff oder Niederalkyl besitzt;

iii) wenn A¹=A²=A³=A⁴ ein Rest der Formel (a-1) oder (a-2) ist und L einen Rest der Formel (b-3)

darstellt, worin X für O steht, Y NR¹⁶, O oder eine direkte Bindung darstellt, und X^a O ist,

a) Y^a dann nicht für steht;

b) und wenn W² Niederalkyl ist, Y^a dann keine direkte Bindung darstellt.

2. Chemische Verbindung nach Anspruch 1, worin L ein Rest der Formel (b-1) ist, worin Y für NH steht,

X O bedeutet und W Wasserstoff ist; oder L ein Rest der Formel (b-1) ist, worin X für S, NH oder NCN steht,

Y NH darstellt und W 1-Piperidinyl, 1-Pyrrolidinyl, 4-Morpholinyl, oder ein Rest der Formel (c-1-c) ist, worin

Z¹ NR¹⁶ darstellt und W¹ Amino, Nitro oder Niederalkyl, gegebenenfalls substituiert mit einem Hydroxy-,

Niederalkylthio, 1-Piperidinyl-, 1-Pyrrolidinyl-, 4-Morpholinyl- oder Phenylrest oder mit zwei

Niederalkylresten, bedeutet; oder

L ein Rest der Formel (b-1) ist, worin X für S, NH oder NCN steht, Y NH darstellt und W ein

Niederalkyloxy oder Niederalkythio ist; oder worin L ein Rest der Formel (b-1) ist, worin W ein Rest der Formel (c-1-a) oder (c-1-b); ist oder

L ein Rest der Formel (b-2) ist, worin n den Wert 1 besitzt, X für O oder S steht und W ein Rest der Formel (c-1-c) ist, worin Z¹ für NR⁶ steht und W¹ Niederalkyl ist; oder

L ein Rest der Formel (b-3) ist, worin X für O steht, Y NH darstellt, X^a O bedeutet, Y^a NR¹⁶ darstellt und W² Niederalkyl ist; oder

L ein Rest der Formel (b-4) ist, worin T ein Rest der Formel (c-3-a) ist, worin X für O oder S steht, Z NR⁸ ist und R⁶ Wasserstoff oder Niederalkyl bedeutet; oder worin T ein Rest der Formel (c-3-b) ist, worin R⁸ Wasserstoff ist und R⁷ Niederalkyl bedeutet; oder

L ein Rest der Formel (b-5) ist, worin Het ein Rest der Formel (c-4-a) ist, worin R⁹, R¹¹ und R¹² Wasserstoff sind; oder worin Het ein Rest der Formel (c-4-c) ist; oder worin Het ein Furan, substituiert mit Niederalkyl, darstellt, welches mit Hydroxy oder mit einem Rest der Formel (c-4-d-1) substituiert ist, worin Y für O oder S steht, Z NH oder eine direkte Bindung bedeutet und R¹⁴ Wasserstoff ist; oder

L ein Rest der Formel (b-6) ist, worin Y¹ für O steht; ist oder L ein Rest der Formel (b-7) ist, worin Ar¹ mit Hydroxy oder Niederalkyloxy substituiertes Phenyl ist.

3. Chemische Verbindung nach einem der Ansprüche 1 und 2 zur Verwendung als Arzneimittel.

4. Chemische Verbindung nach einem der Ansprüche 1 und 2 zur Verwendung als antiallergisches Arzneimittel.

5. Pharmazeutische Zusammensetzung, umfassend einen inerten Träger und eine pharmazeutisch annehmbare Menge einer Verbindung, nach einem der Ansprüche 1 und 2.

6. Pharmazeutische Zusammensetzung nach Anspruch 5 zur Verwendung als anti-allergisches Arzneimittel.

7. Verfahren zur Herstellung einer pharmazeutischen Zusammensetzung, dadurch gekennzeichnet, daß eine therapeutisch wirksame Menge einer Verbindung nach einem der Ansprüche 1 und 2 mit einem geeigneten pharmazeutischen Träger innig vermischt wird.

8. Verfahren zur Herstellung einer Verbindung nach einem der Ansprüche 1 und 2, gekennzeichnet durch:

a) Alkylieren eines Piperidins der Formel Q²-D (III) mit einer Zwischenverbindung der Formel Q¹ (II) in einem reaktionsinerten Lösungsmittel, worin

1) Q² Wasserstoff ist und Q¹ eine Zwischenverbindung der Formel L—G, (II-a) darstellt, worin G Halogen oder Sulfonyloxy bedeutet; oder

2) Q¹ eine Zwischenverbindung der Formel W—C(=X)—G¹, (II-b-1) ist, worin G¹ Halogen, Sulfonyloxy, Niederalkyloxy, Niederalkythio, Ar²-oxy, Ar²-thio, Niederalkylcarbonyloxy oder Niederalkyloxy-carbonyloxy darstellt, oder wenn G¹ an




gebunden ist, es auch —N(Niederalkyl)NO sein kann; und Q² ein Rest der Formel HY¹—Alk— ist, wodurch eine Verbindung der Formel




hergestellt wird; oder

3) Q¹ eine Zwischenverbindung der Formel



ist, und Q² einen Rest der Formel HY¹—Alk— darstellt, wodurch eine Verbindung der Formel



hergestellt wird; oder

4) Q¹ eine Zwischenverbindung der Formel (Niederalkenyl)-G, (II-b-3), ist, und Q² einen Rest HY¹—Alk— darstellt, wodurch eine Verbindung der Formel (Niederalkenyl)—Y¹—Alk—D (I-a-3) hergestellt wird; oder

5) Q¹ eine Zwischenverbindung der Formel




(II-b-4)

ist, und Q^2 einen Rest der Formel




darstellt, wodurch eine Verbindung der Formel



(I-a-4)

hergestellt wird; oder

6) Q¹ eine Zwischenverbindung der Formel $W-C(=X)-Y^1H$, (II-c-1), ist und Q² einen Rest der Formel $G-Alk$ darstellt, wodurch eine Verbindung der Formel (I-a-1) hergestellt wird;



(II-c-2),

ist und Q^2 einen Rest der Formel G—Alk darstellt, wodurch eine Verbindung der Formel (I-a-2) hergestellt wird:

8) Q^1 eine Zwischenverbindung der Formel (Niederalkenyl)— Y^1 —H, (II-c-3), ist und Q^2 einen Rest der Formel $G-Alk-$ darstellt, wodurch eine Verbindung der Formel (I-a-3) hergestellt wird;

9) Q¹ eine Zwischenverbindung der Formel Het'-H, (II-c-4), ist, worin das genannte Het' einen Rest der Formel (c-4-a), (c-4-b) oder (c-4-c) darstellt und Q² ein Rest der Formel G-Alk ist, wodurch eine Verbindung der Formel Het'-Alk-D, (I-a-5), hergestellt wird, oder

b) Umsetzen eines Reagens der Formel W^1-Z^1H , (II-d), mit einer Zwischenverbindung der Formel $HY^1-Alk-D$, (III-b-1), in Gegenwart eines geeigneten $C=X$ bildenden Mittels in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel $W^1-Z^1-C(=X)-Y^1-Alk-D$, (I-a-1-a); oder

c) Umsetzen eines Reagens der Formel (II-d) mit einer Zwischenverbindung der Formel




(II-b-2)

in Gegenwart eines geeigneten




bildenden Mittels, in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel



(I-a-4-a)


hergestellt wird; oder

d) Cyclodesulfurieren einer Zwischenverbindung der Formel




mit einem geeigneten Alkylhalogenid, Metalloxid oder Metallsalz in einem reaktionsinerten Lösungsmittel; oder

10 e) Umsetzen einer Zwischerverbindung der Formel W^1-Z^1-H , (V), mit einem Piperidin der Formel $X^1=C=N-Alk-D$, (VI), worin X^1 für O oder S steht, in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel



hergestellt wird; oder

f) Umsetzen einer Zwischenverbindung der Formel $W^1-N=C-X^1$, (VII) mit einem Piperidin der Formel $HY^1-Alk-D$, (III-b-1), in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel




hergestellt wird; oder

g) Umsetzen einer Zwischenverbindung der Formel (VII) mit einem Piperidin der Formel



30 in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel




40 hergestellt wird; oder

h) Umsetzen einer Zwischenverbindung der Formel $W^2-C(=X^1)-OH$, (VIII), worin W^2 die früher beschriebene Bedeutung von W besitzt, mit der Maßgabe, daß W^2 eine andere Bedeutung als die eines Restes der Formel (c-1-c) aufweist, mit einem Piperidin der Formel $HY^1-Alk-D$ (III-b-1) in einem reaktionsinerten Lösungsmittel, wenn gewünscht, nach Überführen der OH-Funktion in (VIII) in eine geeignete Leaving-Gruppe, oder, wenn gewünscht, durch Umsetzen von (III-b-1) mit (VIII) gemeinsam mit einem entsprechenden Reagens, welches fähig ist, Amide oder Ester zu bilden, wodurch eine Verbindung der Formel




hergestellt wird; oder

i) Umsetzen einer Zwischenverbindung der Formel $W^2-C(=X^1)-OH$, (VIII), worin W^2 die früher definierte Bedeutung besitzt, mit einem Piperidin der Formel



60 in einem reaktionsinerten Lösungsmittel, wenn gewünscht, nach Überführen der OH-Funktion in (VIII) in eine geeignete Leaving-Gruppe, oder, wenn gewünscht, durch Umsetzen von (III-b-2) mit (VIII), gemeinsam mit einem entsprechenden Reagens, das fähig ist, Amide oder Ester zu bilden, wodurch eine Verbindung der Formel



(I-c-2); or


hergestellt wird; oder

j) Umsetzen eines Piperidins der Formel HD, (III-a), mit einem Reagens der Formel L¹-Niederalkendiyli-H, (IX), worin L¹ so ausgewählt ist, daß es gemeinsam mit —Alk— einen Rest der Formel (b-1), (b-3), (b-4), (b-5), (b-6) oder (b-7) bildet, in einem geeigneten reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel L¹—Alk—D, (I-d), hergestellt wird; oder

k) Umsetzen eines Furans der Formel




mit einem Piperidin der Formel (III-a) in Gegenwart von Formaldehyd oder einer polymeren Form hiervon in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel



(I-e)

worin R¹⁸ ein wie in Anspruch 1 definierter Substituent des genannten Furanringes ist, hergestellt wird; oder

l) Alkylieren eines Furanderivates der Formel H—Y¹—D¹, (X-a), worin D¹ einen Rest der Formel



darstellt, mit einer Zwischenverbindung der Formel E—Z—C₆H₂₅—G, (XI-a), worin E einen Rest der Formel



darstellt, in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel E—Z—C₆H₂₅—Y¹—D¹, (I-e-1), hergestellt wird;

m) Alkylieren einer Zwischenverbindung der Formel E—Z—C₆H₂₅—Y¹H, (XI-b), mit einem Furanderivat der Formel G—D¹, (X-b), in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel (I-e-1) hergestellt wird;

n) Alkylieren eines Furanderivates der Formel H—Z¹—C₆H₂₅—Y—D¹, (X-c), mit einer Zwischenverbindung der Formel E—G, (XI-c), in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel




(I-e-2),

hergestellt wird;

o) Alkylieren einer Zwischenverbindung der Formel E—Z¹H, (XI-d), mit einem Furanderivat der Formel G—C₆H₂₅—Y—D¹, (X-d), in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel (I-e-2) hergestellt wird;


p) Umsetzen einer Zwischenverbindung der Formel



(XII-a),

worin Y^{a-1} die früher beschriebene Bedeutung von Y^a besitzt, mit der Maßgabe, daß es keine direkte Bindung darstellt, mit einem Reagens der Formel W^a—C(=X^a)—G¹, (XIII-a) in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel

EP 0 151 824 B1




hergestellt wird;

q) Umsetzen einer Zwischenverbindung der Formel



mit einem Reagens der Formel $\text{R}^6-\text{Z}-\text{C}(=\text{X})-\text{G}^1$ in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel



hergestellt wird;

r) Umsetzen einer Zwischenverbindung der Formel




mit einem Reagens der Formel $\text{R}^7-\text{SO}_2-\text{G}^1$ in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel




hergestellt wird;

s) Umsetzen eines Reagens der Formel $\text{R}^6-\text{N}=\text{C}=\text{X}^{a-1}$, (XIV-a) worin X^{a-1} für O oder S steht, mit einer Zwischenverbindung der Formel (XII-a) in einem reaktionsinerten Lösungsmittel, wodurch eine Verbindung der Formel




hergestellt wird;

t) Umsetzen einer Zwischenverbindung der Formel (XII-b) mit einem Reagens der Formel $\text{R}^6-\text{N}=\text{C}=\text{X}^1$, (XIV-b), wodurch eine Verbindung der Formel



hergestellt wird,


worin D einen Rest der Formel



darstellt; oder
gegebenenfalls Überführen der Verbindungen der Formel (I) ineinander unter Verwendung
techniken bekannter Umwandlungsverfahren für funktionelle Gruppen; und, wenn gewünscht, Überführen
der Verbindungen der Formel (I) in eine therapeutisch wirksame, nicht-toxische Säureadditionssalzform
durch Behandlung mit einer geeigneten Säure oder umgekehrt, Überführen des Säureadditionssalzes mit
Alkali in die freie Basenform; und/oder Herstellen stereochemisch-isomerer Formen hiervon.

Revendications

1. Composé chimique ayant la formule



(I),

sel d'addition acide pharmaceutiquement acceptable ou forme stéréochimiquement isomère possible de
ce dernier, dans lequel:

$A^1=A^2-A^3=A^4$ est un radical bivalent ayant la formule




où un ou deux atomes d'hydrogène dans lesdits radicaux (a-1) — (a-5), peuvent, chacun indépendamment
l'un de l'autre, être remplacés par un halo, un alkyle inférieur, un alkyloxy inférieur, un trifluorométhyle ou
un hydroxy;


R est un membre choisi dans le groupe consistant en hydrogène et alkyle inférieur;

R¹ est un membre choisi dans le groupe consistant en hydrogène, alkyle, cycloalkyle, Ar¹ et alkyle
inférieur substitués par un ou deux radicaux Ar¹;

R² est un membre choisi dans le groupe consistant en hydrogène, alkyle inférieur, cycloalkyle, (alkyle
inférieur)—CO—, (alkyloxy inférieur)—CO— et Ar²-alkyle inférieur; et

L est un radical de formule





(b-4) ;

Het-Alk-

(b-5) ;

(alkényle inférieur)-Y¹-Alk-


(b-6) ; ou

- i) où A¹=A²=A³=A⁴ est un radical de formule (a-3), (a-4) ou (a-5), ou
 ii) où A¹=A²=A³=A⁴ est un radical de formule (a-1) ou (a-2), et R¹ est Ar³ ou un alkyle inférieur substitué par un ou deux radicaux Ar³, ledit Ar³ étant un pyrazinyle, un thiazolyle ou un imidazolyle, éventuellement substitué par un alkyle inférieur ;
 L peut également être un radical de formule


Ar¹-Alk-

(b-7) ;

ledit W étant un membre choisi dans le groupe consistant en hydrogène, alkyle inférieur, Ar¹, Ar¹-alkyle inférieur, 1-pipéridinyle, 1-pyrrolidinyle, 4-morpholinyle, un radical de formule



(c-1-a) ,



(c-1-b) , ou

un radical de formule

un radical de formule

W¹-Z¹-

(c-1-c),

- où R³ et R⁴ sont chacun indépendamment un hydrogène ou un alkyle inférieur ; et W¹ est un cycloalkyle ou un alkyle inférieur, substitués éventuellement par jusqu'à deux substituants choisis dans le groupe consistant en hydroxy


alkyloxy inférieur, 1-pipéridinyle, 1-pyrrolidinyle, 4-morpholinyle et Ar¹ ; et où Z¹ est NR³, W¹ peut aussi être un hydrogène, un amino, un alkylamino inférieur, un Ar¹-amino ou un nitro ;

ledit W² étant un membre choisi dans le groupe consistant en hydrogène, alkyle inférieur, Ar¹ et un radical de formule


R⁶-Z¹-

(c-2-a),

- où R⁶ est un hydrogène, un alkyle inférieur ou Ar¹ ; ledit T étant un radical de formule




- R⁶ étant un hydrogène, un alkyle inférieur ou Ar¹ ;
 R⁷ étant un alkyle inférieur ou Ar¹ ; et
 R⁸ étant un hydrogène ou un alkyle inférieur ;
 ledit Het étant un radical de formule (c-1-a), (c-1-b) ou un radical de formule



(c-4-a)

ou de formule




(c-4-b) ,

où R^9 , R^{10} , R^{11} et R^{12} sont chacun indépendamment un hydrogène ou un alkyle inférieur; ou un radical de formule



où R^{13} est un hydrogène, un alkyle inférieur ou un amino, ou ledit Het étant un furane substitué par un alkyle inférieur, ledit alkyle inférieur étant éventuellement substitué par un hydroxy, un mercapto, un alkyloxy inférieur, un alkythio inférieur, (amino alkyle inférieur) thio, Ar^1-O- ou par un radical de formule



s étant un nombre entier de 1 à 6 inclus; ou bien où Z ou Y est une liaison directe, s peut également être O; et R^{14} étant un hydrogène ou un alkyle inférieur;

où n = O ou le nombre entier 1 ou 2;

X est O, S, NR^{15} ou $CHNO_2$;

Y est O, S, NR^{16} ou une liaison directe;

Y^1 est O, S ou NR^{16} ;

Y^2 est S ou NR^{16} ;

Z est O, S, NR^8 ou une liaison directe;

Z^1 est O, S ou NR^8 ;

X^8 et Y^8 ayant indépendamment respectivement la même valeur que X et Y;

ledit R^{15} étant un hydrogène, un alkyle inférieur, cyano, nitro, Ar^2 -sulfonyl, alkylsulfonyl inférieur, alkylcarbonyl inférieur ou Ar^2 -carbonyl;

ledit R^{16} étant un hydrogène, alkyle inférieur (Ar^3)alkyle inférieur, 2-alkyloxy inférieur-1,2-dioxéthyle; ou un radical de formule $-C(=X)-R^{17}$; R^{17} étant un hydrogène, alkyle inférieur, Ar^2 -alkyle inférieur, alkyloxy inférieur, Ar^2 -alkyloxy inférieur, mono- ou di(alkyle inférieur)amino, Ar^2 -alkylamino inférieur ou Ar^2 -alkyle inférieur(alkyle inférieur) amino;

où Ar^1 est un membre choisi dans le groupe consistant en phényle, étant éventuellement substitué par jusqu'à trois substituants, chacun choisi indépendamment dans le groupe consistant en halo, hydroxy, nitro, cyano, trifluorométhyle, alkyle inférieur, alkyloxy inférieur, alkythio inférieur, mercapto, amino, mono- et di(alkyle inférieur) amino, carboxyle, alkylcarbonyl inférieur et alkyle inférieur-CO—; thiényl; halothiényl; furanyl; furanyl substitué par un alkyle inférieur; pyridinyl; pyrazinyl; thiazolyl et imidazolyl substitué éventuellement par un alkyle inférieur; et où Ar^2 est un membre choisi dans le groupe consistant en phényle substitué éventuellement par jusqu'à trois substituants chacun choisi indépendamment dans le groupe consistant en halo, hydroxy, nitro, cyano, trifluorométhyle, alkyle inférieur, alkyloxy inférieur, alkythio inférieur, mercapto, amino, mono- et di(alkyle inférieur)amino, carboxyle, alkylcarbonyl inférieur et (alkyle inférieur)-CO; où l'alkyle inférieur représente des radicaux hydrocarbures saturés à chaînes droites et ramifiées ayant 1 à 6 atomes de carbone; l'alkyle

comprend un alkyle inférieur et l'homologue supérieur de ce dernier ayant 7 à 10 atomes de carbone; le cycloalkyle est un cyclopropyle, cyclobutyle, cyclopentyle et cyclohexyle; l'alkényle inférieur représente un radical hydrocarbure à chaîne droite ou ramifiée contenant une double liaison et ayant 2 à 6 atomes de carbone; l'alkanédiyle inférieur représente des radicaux alkanédiyle bivalents à chaînes droites ou ramifiées ayant de 1 à 6 atomes de carbone;

à condition que:

i) lorsque $A^1=A^2=A^3=A^4$ est un radical de formule (a-1) ou (a-2), et L est un radical de formule (b-1), où W est autre qu'un hydrogène ou autre qu'un radical de formule (c-1-a) ou (c-1-b), alors X est différent de O;

ii) lorsque L est un radical de formule (b-1), où W est un radical de formule (c-1-c), où Z^1 est NH, alors W^1 est autre qu'un hydrogène ou un alkyle inférieur;

iii) lorsque $A^1=A^2=A^3=A^4$ est un radical de formule (a-1) ou (a-2), et L est un radical de formule (b-3), où X est O, Y est NR^{15} , O ou une liaison directe, et X^8 est O,

a) alors Y^8 n'est pas O;

b) et W^2 étant un alkyle inférieur, alors Y^8 n'est pas une liaison directe.

2. Composés chimiques selon la revendication 1, dans lequel L est un radical de formule (b-1), dans lequel Y est NH, X et O et W est un hydrogène; ou L est un radical de formule (b-1) dans lequel X est S, NH ou CN, Y est NH et W est un 1-pipéridinyle, 1-pyrrolidinyle, 4-morpholinyle, ou un radical de formule (c-1-c), dans lequel Z^1 est NR^8 et W^1 est un amino, nitro, alkyle inférieur, éventuellement substitué par un radical hydroxy, alkyloxy inférieur, 1-pipéridinyle, 1-pyrrolidinyle, 4-morpholinyle ou phényle, ou par deux radicaux alkyloxy inférieur; ou L est un radical de formule (b-1), dans lequel X est S, NH ou CN, Y est NH et W est un alkyloxy inférieur ou un alkythio inférieur; ou dans lequel L est un radical de formule (b-1) où W

est un radical de formule (c-1-a) ou (c-1-b); ou

L est un radical de formule (b-2) où n est 1, X est O ou S et W est un radical de formule (c-1-c) où Z¹ est NR⁸ et W¹ est un alkyle inférieur; ou

L est un radical de formule (b-3) où X est O, Y est NH, X^a est O, Y^a est NR¹⁶ et W² est un alkyle inférieur;

ou

L est un radical de formule (b-4) où T est un radical de formule (c-3-a) où X est O ou S, Z est NR⁹ et R⁶ est un hydrogène ou un alkyle inférieur; ou dans lequel T est un radical de formule (c-3-b) où R⁸ est un hydrogène et R⁷ est un alkyle inférieur; ou

L est un radical de formule (b-5) où Het est un radical de formule (c-4-a) où R⁹, R¹¹ et R¹² sont des hydrogènes; ou encore où Het est un radical de formule (c-4-c); ou encore où Het est un furane substitué par un alkyle inférieur étant substitué par un hydroxy ou par un radical de formule (c-4-d-1), où Y est O ou S, Z est NH ou une liaison directe et R¹⁴ est un hydrogène; ou

L est un radical de formule (b-6) où Y¹ est O; ou

L est un radical de formule (b-7) où Ar¹ est un phényle substitué par un hydroxy ou un alkyloxy inférieur.

3. Composé chimique selon l'une quelconque des revendications 1 et 2 pour être utilisé comme médicament.

4. Composé chimique selon l'une quelconque des revendications 1 et 2 pour être utilisé comme médicament anti-allergique.

5. Composition pharmaceutique comprenant un excipient inerte et une quantité pharmaceutiquement acceptable de composé selon l'une quelconque des revendications 1 et 2.

6. Composition pharmaceutique selon la revendication 5 pour être utilisée comme médicament anti-allergique.

7. Procédé pour la préparation d'une composition pharmaceutique, caractérisé en ce que une quantité thérapeutiquement efficace de composé selon l'une quelconque des revendications 1 et 2 est intimement mélangée avec un excipient pharmaceutique approprié.

8. Procédé pour la préparation d'un composé selon l'une quelconque des revendications 1 et 2, caractérisé de la façon suivante:

a) en alkylant une pipéridine de formule Q²—D (II) avec un intermédiaire de formule Q¹ (II) dans un solvant inerte, où

1) Q² est un hydrogène et Q¹ est un intermédiaire de formule L—G (II-a), ledit G représentant un halo ou un sulfonyloxy; ou


2) Q¹ est un intermédiaire de formule W—C(=X)—G¹, (II-b-1), ledit G¹ représentant un halo, sulfonyloxy, alkyloxy inférieur, alkylthio inférieur, Ar²-oxy, Ar²-thio, alkylcarbonyloxy inférieur ou alkylxycarbonyloxy inférieur, ou encore où G¹ est relié à




il peut également être un —N(alkyle inférieur)NO; et Q² est un radical de formule HY¹—Alk—, préparant ainsi un composé de formule



3) Q¹ est un intermédiaire de formule



et Q² est un radical de formule HY¹—Alk—, préparant ainsi un composé de formule



4) Q¹ est un intermédiaire de formule (alkényle inférieur)—G, (II-b-3), et Q² est un radical HY¹—Alk— préparant ainsi un composé de formule (alkényle inférieur)—Y¹—Alk—D (I-a-3); ou

5) Q^1 est un intermédiaire de formule




(11-b-4)

et Q^2 est un radical de formule




préparant ainsi un composé de formule



(I-a-4); ou

6) Q¹ est un intermédiaire de formule W—C(=X)—Y¹H (II-c-1) et Q² est un radical de formule G—Alk, préparant ainsi un composé de formule (I-a-1);
7) Q¹ est un intermédiaire de formule



(II-c-2),

et Q² est un radical de formule G—Alk, préparant ainsi un composé de formule (I-a-2);

8) Q¹ est un intermédiaire de formule (alkényle inférieur)—Y—H (II-c-3), et Q² est un radical de formule G—Alk—, préparant ainsi un composé de formule (I-a-3);

9) Q¹ est un intermédiaire de formule Het'—H, (II-c-4), où ledit Het' est un radical de formule (c-4-a), (c-4-b) ou (c-4-c) et Q² est un radical de formule G—Alk, préparant ainsi un composé de formule Het'—Alk—D (I-a-5); ou

b) en faisant réagir un réactif de formule W^1-Z^1-H (II-d) avec un intermédiaire de formule $HY^1-Alk-D$ (III-b-1) en présence d'un agent approprié générant $C=X$, dans un solvant inerte, préparant ainsi un composé de formule $W^1-Z^1-C(X)-Y^1-Alk-D$, (I-a-1-a); ou




(II-b-2)

en présence d'un agent approprié générant




dans un solvant inerte, préparant ainsi un composé de formule



(I-a-4-a); ou


d) en cyclodésulfurant un intermédiaire de formule



(IV)

EP 0 151 824 B1

avec un halogénure d'alkyle, un oxyde métallique ou un sel métallique approprié dans un solvant inerte; ou
e) en faisant réagir un intermédiaire de formule W^1-Z^1-H (V) avec une pipéridine de formule $X^1=C=N-Alk-D$ (VI), où X^1 est O ou S, dans un solvant inerte, préparant ainsi un composé de formule



f) en faisant réagir un intermédiaire de formule $W^1-N=C=X^1$ (VII) avec une pipéridine de formule $HY^1-Alk-D$ (III-b-1), dans un solvant inerte, préparant ainsi un composé de formule



g) en faisant réagir un intermédiaire de formule (VII) avec une pipéridine de formule



dans un solvant inerte, préparant ainsi un composé de formule



h) en faisant réagir un intermédiaire de formule $W^2-C(=X^1)-OH$ (VIII), ledit W^2 ayant la valeur décrite précédemment de W , à condition que W^2 soit autre qu'un radical de formule (c-1-c), avec une pipéridine de formule $HY^1-Alk-D$ (III-b-1), dans un solvant inerte, si on le désire, après avoir transformé la fonction OH dans (VIII) en un groupe partant approprié, ou, si on le désire, en faisant réagir (III-b-1) avec (VIII) avec un réactif approprié capable de former des amides ou des esters, préparant ainsi un composé de formule



i) en faisant réagir un intermédiaire de formule $W^2-C(=X^1)-OH$ (VIII), ledit W^2 ayant la valeur définie précédemment, avec une pipéridine de formule



dans un solvant inerte, si on le désire, après avoir transformé la fonction OH dans (VIII) en un groupe partant approprié, ou, si on le désire, en faisant réagir (III-b-2) avec (VIII), avec un solvant approprié capable de former des amides ou des esters, préparant ainsi un composé de formule




j) en faisant réagir une pipéridine de formule HD (III-a) avec un réactif de formule L^1 -alkényldiyle inférieur-H (IX), où L^1 est choisi de telle sorte qu'il forme, combiné avec $-Alk-$, un radical de formule (b-1), (b-3), (b-4), (b-5), (b-6) ou (b-7), dans un solvant inerte approprié, préparant ainsi un composé de formule $L^1-Alk-D$ (I-d); ou

k) en faisant réagir un furane de formule




EP 0 151 824 B1

avec une pipéridine de formule (III-a) en présence de formaldéhyde ou d'une forme polymère de celui-ci, dans un solvant inerte, préparant ainsi un composé de formule



où R^{1a} est un substituant, défini dans la revendication 1, dudit cycle furane; ou


10 l) en alkylant un dérivé du furane de formule H-Y¹-D¹, (X-a), où D¹ représente un radical de formule



avec un intermédiaire de formule E-Z-C₆H_{2a}-G (XI-a), où E représente un radical de formule



dans un solvant inerte, préparant ainsi un composé de formule




m) en alkylant un intermédiaire de formule E-Z-C₆H_{2a}-Y¹H, (XI-b), avec un dérivé du furane de formule G-D¹, (X-b), dans un solvant inerte, préparant ainsi un composé de formule (I-e-1);

30 n) en alkylant un dérivé du furane de formule H-Z¹-C₆H_{2a}-Y-D¹, (X-c), avec un intermédiaire de formule E-G, (XI-c), dans un solvant inerte, préparant ainsi un composé de formule




35 o) en alkylant un intermédiaire de formule E-Z¹-H (XI-d), avec un dérivé du furane de formule G-C₆H_{2a}-Y-D¹, (X-d), dans un solvant inerte, préparant ainsi un composé de formule (I-e-2);


p) en faisant réagir un intermédiaire de formule




45 où Y^{a-1} a été décrit précédemment comme ayant la valeur de Y^a, à condition qu'il ne soit pas une liaison directe, avec un réactif de formule W²-C(=X^a)-G¹, (XII-a), dans un solvant inerte, préparant ainsi un composé de formule



q) en faisant réagir un intermédiaire de formule




avec un réactif de formule R⁶-Z-C(=X)-G¹, dans un solvant inerte, préparant ainsi un composé de formule




EP 0 151 824 B1


r) en faisant réagir un intermédiaire de formule




avec un réactif de formule $\text{R}^7 - \text{SO}_2 - \text{G}^1$, dans un solvant inerte, préparant ainsi un composé de formule




15 s) en faisant réagir un réactif de formule $\text{R}^5 - \text{N}=\text{C}=\text{X}^{a-1}$ (XIV-a), où X^{a-1} est O ou S, avec un intermédiaire de formule (XII-a), dans un solvant inerte, préparant ainsi un composé de formule



25 t) en faisant réagir un intermédiaire de formule (XII-b) avec un réactif de formule $\text{R}^6 - \text{N}=\text{C}=\text{X}^1$ (XIV-b), préparant ainsi un composé de formule



où D représente un radical de formule



40 en transformant éventuellement les composés de formule (I) l'un en l'autre conformément aux procédés connus de transformation des groupes fonctionnels; et, si on le désire, en transformant les composés de formule (I) en une forme de sel d'addition acide thérapeutiquement active non toxique par traitement avec un acide approprié ou, inversement, en transformant le sel d'addition acide en la forme de base libre avec un alcali; et/ou en préparant leurs formes stéréochimiquement isomères.

50

55

60

65